

Supplementary Materials for

Microbiological ascertainment in patients with pneumonia: the experience of a teaching hospital in Rome

**Marianxhela Dajko, Andrea Poscia, Brunella Posteraro, Domenico Speciale,
Massimo Volpe, Silvia Mancinelli, Walter Ricciardi and Chiara de Waure**

[Early release]

Corresponding author:

Marianxhela Dajko, Centro di Epidemiologia, Biostatistica e Informatica Medica, Università Politecnica delle Marche, Via Tronto 10/A, 60120 Torrette di Ancona, Ancona, Italy. E-mail: marianxheladajko@hotmail.com.

Published on

Ann Ist Super Sanità 2020 Vol. 56, No. 3
DOI: 10.4415/ANN_20_03

This PDF file includes:

Table. Frequency of the etiological agent

Table
Frequency of the etiological agent

Pathogens	N (%) ^a
Achromobacter xylosoxidans	1 (0.1)
Acinetobacter baumannii	170 (22.9)
Other Acinetobacter	4 (0.5)
Aeromonas caviae	2 (0.3)
Aspergillus	23 (3.1)
Beauveria bassiana	1 (0.1)
Blastoschizomyces capitatus	5 (0.7)
Moraxella catarrhalis	2 (0.3)
Brucella abortus	2 (0.3)
Burkholderia cepacian	1 (0.1)
Candida albicans	278 (37.4)
Other Candida	147 (19.8)
Chaetomium jodhpurens	1 (0.1)
Citrobacter	10 (1.3)
Cladosporium cladosporioides	1 (0.1)
Corynebacterium	2 (0.3)
Enterobacter	23 (3.1)
Enterococcus faecalis tipo D	19 (2.6)
other Enterococcus	19 (2.6)
Escherichia coli	47 (6.3)
Flavobacterium meningoseptum	1 (0.1)
Fusarium oxisporum	2 (0.3)
Fusarium solani	1 (0.1)
Geotrichum candidum, D-galactose	1 (0.1)
Haemophilus influenzae	35 (4.7)
Hafnia alvei	1 (0.1)
Klebsiella pneumoniae	74 (10)
Other Klebsiella	24 (3.2)
Mycobacterium avium complex ("MAC")	3 (0.4)
Mycobacterium tuberculosis	5 (0.7)
Neisseria meningitidis	1 (0.1)
Neisseria sicca	282 (38)
Pichia kluyveri	2 (0.3)
Pneumocystis carinii jiroveci	7 (0.9)
Proteus	21 (2.8)
Providencia stuartii	1 (0.1)
Pseudomonas aeruginosa	102 (13.7)
Pseudomonas putida	3 (0.4)
Rhizomucor pusillus	1 (0.1)
Saccharomyces cerevisiae	9 (1.2)
Serratia marcescens	18 (2.4)
Sphingomonas paucimobilis	1 (0.1)
Staphylococcus aureus	82 (11)
Staphylococcus haemolyticus	15 (2)
other Staphylococcus	19 (2.6)
Stenotrophomonas maltophilia	40 (5.4)
Streptococcus pneumoniae	299 (40.2)
other Streptococcus	9 (1.2)

^a-% are calculated for 743 patients who had a positive result