

Supplementary Materials for Gender and burden differences in family caregivers of patients affected by ten rare diseases

Flavia Chiarotti, Yllka Kodra, Marta De Santis, Maria Bellenghi, Domenica Taruscio,
Alessandra Carè and Marina Petrini

Corresponding author:

Marina Petrini, Centro di Riferimento Medicina di Genere, Istituto Superiore di Sanità, Viale Regina Elena 299, 00161 Rome, Italy. Email: marina.petrini@iss.it.

Published on

Ann Ist Super Sanità 2023 Vol. 59, No. 2: 122-131
DOI: 10.4415/ANN_23_02_05

This PDF file includes:

Table 1aS

Family caregivers' Zarit answers to Item 7 "Do you feel fear for the future?" by family caregivers' sex

Table 1bS

Family caregivers' Zarit answers to Item 10 "Do you believe that taking care of your family member has adversely affected your health?" by family caregivers' sex

Table 2S

Family caregivers: burden by sex, occupation and age group

Table 3S

Family caregivers: EQ5D answers to Item 4 "Do you feel pain and discomfort?" by patients' age

Table 1aSFamily caregivers' Zarit answers to Item 7 "Do you feel fear for the future?" by family caregivers' sex ($p=0.007^{**}$)

| FCs sex | FCs number | "Never" n (%) | "Rarely" n (%) | "Sometimes" n (%) | "Very often" n (%) | "Nearly always" n (%) |
|----------------|-------------------|--------------------------|---------------------------|------------------------------|-------------------------------|----------------------------------|
| M | 53 | 4 (7.6) | 5 (9.4) | 21 (39.6) | 7 (13.2) | 16 (30.2) |
| F | 146 | 2 (1.4) | 8 (5.5) | 36 (24.7) | 43 (29.4) | 57 (39.0) |
| Total | 199 | 6 (3.0) | 13 (6.5) | 57 (28.6) | 50 (25.1) | 73 (36.7) |

p: significance level of the Fisher's exact probability test: ** $p \leq 0.01$; FC: family caregiver; M: males; F: females; n (%): absolute and percent frequency of FC responders each scale grade of the Zarit Item 7.

Table 1bSFamily caregivers' Zarit answers to Item 10 "Do you believe that taking care of your family member has adversely affected your health?" by family caregivers' sex ($p=0.012^*$)

| FCs sex | FCs number | "Never" n (%) | "Rarely" n (%) | "Sometimes" n (%) | "Very often" n (%) | "Nearly always" n (%) |
|----------------|-------------------|--------------------------|---------------------------|------------------------------|-------------------------------|----------------------------------|
| M | 53 | 29 (54.7) | 5 (9.4) | 15 (28.3) | 2 (3.8) | 2 (3.8) |
| F | 146 | 46 (31.5) | 30 (20.5) | 51 (34.9) | 17 (11.6) | 2 (1.4) |
| Total | 199 | 75 (37.7) | 35 (17.6) | 66 (33.2) | 19 (9.5) | 4 (2.0) |

p: significance level of the Fisher's exact probability test: * $p \leq 0.05$; FC: family caregiver; M: males; F: females; n (%): absolute and percent frequency of FC responders each scale grade of the Zarit Item 10.

Table 2S

Family caregivers: burden by sex, occupation and age group

| FCs occupation | Zarit score | Male FCs age groups | | | | Female FCs age groups | | | |
|-----------------------|--------------------|----------------------------|----------------|--------------|--------------|------------------------------|----------------|--------------|--------------|
| | | -39 y | 40-54 y | 55- y | Total | -39 y | 40-54 y | 55- y | Total |
| Occupied | n (%) | 6 (85.7) | 26 (86.7) | 7 (46.7) | 39 (75.0) | 28 (58.3) | 42 (52.5) | 3 (17.6) | 73 (50.3) |
| | mean (SD) | 22.8 (6.9) | 28.5 (14.1) | 22.3 (16.6) | 26.5 (13.7) | 29.1 (12.7) | 25.5 (11.4) | 22.3 (17.1) | 26.8 (12.1) |
| | median (min-max) | 21.5 (14-32) | 29 (10-60) | 22 (0-46) | 22 (0-60) | 28 (6-56) | 25 (7-59) | 14 (11-42) | 25 (6-59) |
| Retired | n (%) | 0 (0) | 0 (0) | 8 (53.3) | 8 (15.4) | 1 (2.1) | 7 (8.8) | 8 (47.1) | 16 (11.0) |
| | mean (SD) | | | 25.0 (10.4) | 25.0 (10.4) | 16.0 (0) | 34.7 (15.3) | 27.9 (11.2) | 30.1 (13.3) |
| | median (min-max) | | | 22 (13-41) | 22 (13-41) | 16 (16-16) | 31 (20-59) | 29 (12-48) | 28 (12-59) |
| Homemaker | n (%) | 0 (0) | 2 (6.7) | 0 (0) | 2 (3.8) | 10 (20.8) | 26 (32.5) | 6 (35.3) | 42 (29.0) |
| | mean (SD) | | 15.0 (1.4) | | 15.0 (1.4) | 24.7 (10.1) | 30.6 (13.3) | 22.5 (11.5) | 28.0 (12.6) |
| | median (min-max) | | 15 (14-16) | | 15 (14-16) | 26 (7-42) | 29 (8-56) | 21 (9-44) | 27 (7-56) |
| Others | n (%) | 1 (14.3) | 2 (6.7) | 0 (0) | 3 (5.8) | 9 (18.8) | 5 (6.3) | 0 (0) | 14 (9.7) |
| | mean (SD) | 21.0 (0) | 27.0 (4.2) | | 25.0 (4.6) | 27.4 (10.2) | 42.4 (10.9) | | 32.8 (12.5) |
| | median (min-max) | 21 (21-21) | 27 (24-30) | | 24 (21-30) | 27 (16-46) | 45 (29-56) | | 31 (16-56) |
| Total | n (%) | 7 (100) | 30 (100) | 15 (100) | 52 (100) | 48 (100) | 80 (100) | 17 (100) | 145 (100) |
| | mean (SD) | 22.6 (6.4) | 27.5 (13.6) | 23.7 (13.2) | 25.8 (12.7) | 27.6 (11.7) | 29 (13) | 25 (11.9) | 28.1 (12.4) |
| | median (min-max) | 21 (14-32) | 26 (10-60) | 22 (0-46) | 22 (0-60) | 27 (6-56) | 27 (7-59) | 21 (9-48) | 27 (6-59) |

FC: family caregiver; y: years; Zarit: burden; others: never occupied or forced to leave work.

SD: standard deviation; min-max: minimum-maximum.

Table 3SFamily caregivers: EQ5D answers to Item 4 "Do you feel pain and discomfort?" by patients' age ($p=0.003^{**}$)

| FCs caring for | FCs number | "No pain or discomfort" n (%) | "Mild pain or discomfort" n (%) | "Moderate pain or discomfort" n (%) | "Severe pain or discomfort" n (%) | "Extreme pain or discomfort" n (%) | Median category |
|-----------------------|-------------------|--|--|--|--|---|---------------------------|
| Children | 137 | 85 (62.0) | 31 (22.6) | 19 (13.9) | 1 (0.7) | 1 (0.7) | "No pain or discomfort" |
| Adults | 56 | 20 (35.7) | 23 (41.1) | 9 (16.1) | 3 (5.4) | 1 (1.8) | "Mild pain or discomfort" |
| Total | 193 | 105 (54.4) | 54 (28.0) | 28 (14.5) | 4 (2.1) | 2 (1.0) | |

FC: Family caregiver.

n (%): absolute and percent frequency of FC responders to each scale grade of the EQ-5D Item 4.

p: significance level of the chi-squared test comparing FCs of child vs adult subjects with respect to Item 4: ** $p \leq 0.01$. Median category (of an ordinal variable): the category whose cumulative frequency is greater than (and nearest to) 50%.