



EXECUTIVE SUMMARY for 2018

IARTR THE ITALIAN ASSISTED REPRODUCTIVE TECHNOLOGY REGISTER

**MONITORING THE ACTIVITY AND
OUTCOMES OF ITALIAN ASSISTED
REPRODUCTIVE TECHNOLOGY CENTERS
2018**

MONITORING THE ACTIVITY AND OUTCOMES OF ITALIAN ART CENTERS IN 2018

AUTHORS

**G. Scaravelli, V. Vigilano, R. De Luca, S. Bolli, R. Spoletini,
A. Bertini, L. Speziale**

*ART Italian National Register, National Centre for Diseases
Prevention and Health Promotion, National Health Institute,
Rome, Italy*

INDEX

SUMMARY OF OUTPUTS GENERATED FROM IARTR, 2018.....	1
THE ITALIAN ASSISTED REPRODUCTION TECHNOLOGY REGISTER (IARTR)	2
HOW DOES IARTR WORK?	3
THE IARTR WEBSITE.....	4
1. ACCESS AND UTILIZATION OF ART SERVICES IN ITALY, 2018	5
1.1. ACCESS TO ART SERVICE.....	6
1.2. UTILIZATION OF ART SERVICES	8
2. EFFICACY OF ART IN 2018 AND TIME TRENDS FROM 2005 TO 2018	12
2.1. OVERVIEW OF ART.....	13
2.1.1. <i>What types of ART cycles were performed?</i>	13
2.1.2. <i>How did the types of ART techniques change according to transfers among women of different ages?</i>	15
2.2. ART, NON-DONOR CYCLES.....	16
2.2.1. <i>What are the causes of infertility of couples in ART treatment using fresh cycles in 2018?</i>	16
2.2.2. <i>What are the steps for an ART treatment using fresh cycles?</i>	17
2.2.3. <i>Did the number of embryos transferred differ among women of different age groups?</i>	18
2.2.4. <i>What are the percentages of initiated cycles or thawings, and transfers that result in pregnancies for ART cycles?</i>	19
2.2.5. <i>What is the “Cumulative Pregnancy Rate”?</i>	20
2.2.6. <i>What percentage of ART pregnancies resulted in a delivery?</i>	21
2.2.7. <i>What is the gender distribution of infertility factors among ART users?</i>	22
2.2.8. <i>Is the use of ART procedures change over time?</i>	23
2.2.9. <i>Did the use of FET (Frozen Embryo Transfer) procedures differ in Italy compared to the other European countries over time?</i>	24
2.2.10. <i>Has the age of ART female patients changed over time?</i>	25

2.2.11. Has the number of embryos transferred changed in fresh cycles?.....	26
2.2.12. Did pregnancy rates per transfer changed over time among different ART procedures?	27
2.2.13. Did cumulative pregnancy rates per initiated cycle changed over time?	28
2.2.14. Does the risk of pregnancy loss differ among women of different age groups?	29
2.3. ART DONOR CYCLES.....	30
2.3.1. Which gametes were used in ART donor cycles in 2018?	30
2.3.2. What was the age of recipient female patients in ART donor cycles in 2018.	31
2.3.3. What was the percentage of transfers that result in pregnancies in ART donor cycles in 2018?	32
2.3.3. What were the outcomes of pregnancies obtained in ART donor cycles in 2018?	33
2.4. PGT (PRE-IMPLANTATION GENETIC TESTING) ACTIVITY	34
2.4.1. Which types of genetic tests were performed in ART cycles in 2018?	34
2.4.2. Did the use of different genetic analysis in ART change over time?	35
3. ART SAFETY INDICATORS	36
3.1. SAFETY IN ART PROCEDURES	37
3.1.1. Did the numbers of complications for ART cycles change over time?	37
3.1.2. Did the percentages of multiple deliveries for ART non donor cycles change over time?	38
3.1.2. Did the percentages of preterm live babies change over time?	40
3.1.4. Did the percentage of underweight live babies change over time?.....	41
4. INTRA-UTERINE INSEMINATION PROCEDURES	42
4.1. ACCESS TO INTRA-UTERINE INSEMINATION SERVICE.....	43
4.2. EFFICACY SAFETY AND TRENDS OF IUI	45
4.2.1. Is the use of IUI-H increasing?	45
4.2.2. Do percentages of IUI-H cycles resulting in pregnancies differ among women of different age groups?.....	46
4.2.3. What were the outcomes of pregnancies obtained in IUI-H cycles in 2018?.	47
4.2.4. Did the numbers of complications for H-IUI cycles change over time?	48

4.2.5. <i>Did the percentages of multiple deliveries change over time for homologous intrauterine insemination cycles?</i>	49
4.3. IUI DONOR CYCLES.	50
4.3.1. <i>What was the outcome in IUI donor cycles in 2018?</i>	50
APPENDIX	51
SUMMARY TABLE OF ACTIVITY AND OUTCOME OF ART PROCEDURES YEARS 2011-2018	51
SUMMARY TABLE OF ACTIVITY AND OUTCOME OF ALL ART PROCEDURES, 2011-2018.....	52
SUMMARY TABLE OF ACTIVITY AND OUTCOME OF ART NON DONOR PROCEDURES, 2011-2018	53
SUMMARY TABLE OF ACTIVITY AND OUTCOME OF ART WITH PGT ANALYSIS, 2014-2018	54
SUMMARY TABLE OF ACTIVITY AND OUTCOME OF ART PROCEDURES WITH GAMETE DONATION, 2014-2018.....	55
BIBLIOGRAPHY	56

Downloadable at: <http://www.iss.it/rpma>

SUMMARY OF OUTPUTS GENERATED FROM IARTR, 2018

	Non-donor Procedures			Gametes donation		
	Fresh cycles (IVF and ICSI)	FET	FO	sperm donation	oocyte donation	double donation
n° of patients	42.090	16.067	1.201	1.187	5.160	378
n° initiated cycles	51.086	-	-	1.340	5.981	422
n° aspirations/thawings	46.387	19.587	1.318	-	-	-
n° transfers	30.584	19.016	1.036	1.063	5.550	397
<i>with 1 embryo</i>	<i>11.955</i>	<i>13.357</i>	<i>364</i>	<i>523</i>	<i>3.166</i>	<i>233</i>
<i>with 2 embryos</i>	<i>15.118</i>	<i>5.207</i>	<i>584</i>	<i>489</i>	<i>2.268</i>	<i>153</i>
<i>with 3 embryos</i>	<i>3.322</i>	<i>434</i>	<i>87</i>	<i>51</i>	<i>116</i>	<i>11</i>
<i>with 4 o + embryos</i>	<i>189</i>	<i>18</i>	<i>1</i>	<i>-</i>	<i>-</i>	<i>-</i>
n° clinical pregnancies*	8.307	5.995	223	401	1.954	162
pregnancies per initiated cycles (%)	16,3	-	-	29,9	32,7	38,4
pregnancies per initiated cycles without freeze-all cycles (%)	19,0	-	-	-	-	-
pregnancies per aspirations/thawings (%)	17,9	30,6	16,9	-	-	-
pregnancies per aspirations/thawings without freeze-all cycles (%)	21,3	-	-	-	-	-
pregnancies per transfers (%)	27,2	31,5	21,5	37,7	35,2	40,8
pregnancies lost to follow-up (%)	10,9	6,6	6,7	11,0	14,0	17,9
n° pregnancy losses	1.943	1.474	55	82	371	28
pregnancies loss per monitored pregnancy (%)	26,3	26,3	26,4	23,0	22,1	21,1
n° deliveries	5.458	4.127	153	275	1.310	105
deliveries per aspirations/thawings (%)	11,8	21,1	11,6	20,5	21,9	24,9
estimated deliveries per aspirations/thawings (%)	13,2	22,6	12,4	23,1	25,5	30,3
estimated deliveries per aspiration without freeze-all cycles (%)	15,7					
twin deliveries (%)	13,6	6,7	12,4	13,5	12,4	12,4
triplets or more deliveries (%)	0,3	0,1	0,0	0,4	0,2	0
n° live born babies	6.186	4.393	172	311	1.466	118

*Clinical pregnancy: A pregnancy diagnosed by ultrasonographic visualization of one or more gestational sacs or definitive clinical signs of pregnancy. In addition to intra-uterine pregnancy, it includes a clinically documented ectopic pregnancy¹.

THE ITALIAN ASSISTED REPRODUCTION TECHNOLOGY REGISTER (IARTR)

IARTR has been established at the *Istituto Superiore di Sanità* (Italian National Institute of Health), *National Centre for Epidemiology Surveillance and Health Promotion*, by a Decree of the Ministry of Health issued on the 7th of October 2005 (G.U. n. 282 del 3rd December 2005) in implementation of article n° 11 paragraph 1 of Law 40/2004 (G.U. n. 45 del 24th February 2004).

The Register collects descriptive, technical, structural and organizational information of Assisted Reproductive Techniques (ART) centers authorized by their regional health authority, to conduct assisted reproductive technology, and anonymous, aggregate data sets on all the ART treatments, plus information on the infertile couples, on embryos created and on children born after ART.

The main objectives of the Register are:

- ASSESS and REGISTER all the centres performing ART treatments and IntraUterine Insemination (IUI) procedures in the country and the number of embryos created and cryopreserved;
- COLLECT and EVALUATE data regarding centres characteristics and addresses, type of service offered (public, private or private covered by the National Health Service), the different techniques performed, activity, availability, efficacy and safety of techniques application;
- PROMOTE research and study on couple infertility causes, long-term evaluation of well-being of the children born after ART procedures; research on gametes characteristics and new cryopreservation protocols; MONITORING time trends in ART applications in order to compare different attitudes with other countries.

The Register prepares an annual epidemiological/statistical report on the ART centres' activity for the Minister of Health in order to illustrate to the Parliament the situation in the ART field with a particular epidemiological overview.

The IARTR is linked to the European IVF Monitoring (EIM – European *In vitro* fertilization Monitoring) Consortium, which collects data on ART from about 39 European countries. In turn, the EIM sends data to the World Register ICMART (International Committee Monitoring Assisted Reproductive Technologies). Professor Jacques de Mouzon, Secretary of the ART World Register (ICMART), audited the activity of IARTR since 2018.

HOW DOES IARTR WORK?

Dr Giulia Scaravelli, MD-Gynaecologist and PhD-Obstetrics and Gynaecology, coordinates the staff. In the staff, there are varieties of skills: statistics, epidemiology, gynaecology, biology, psychology and informatics.

Data on efficacy, safety and outcomes of reproductive techniques, including IUI, are collected on a website on a reserved area with username and password. Data collection is based on summary data sent from each centre according to a national law on privacy protection (Dl. 196/2003).

The data collection is organized in two different periods:

- The first phase of the collection is related to the activity conducted and the results obtained in 2018 and it was carried out from May to June 2019;
- The second phase of the collection is related to the outcomes of pregnancies obtained from ART treatments started in 2018 and it was carried out from October to December 2019.

Data collection is made on number of cycles performed for each technique, number of patients treated, kind of infertility diagnosed, embryos created and eventually transferred, pregnancies outcomes, babies born and complications during treatments.

THE IARTR WEBSITE

www.iss.it/rpma

The Register website has the goal to collect and disseminate data and information related to IUI and ART procedures.

There are different levels of interest in the website:

- A service to the citizens: they can consult the list of all the authorized ART centres by different regions and can have information about the techniques they perform, and the availability of the service. They can find on the home page all the information regarding ART and IUI techniques and their application in Italy. They can find also the links to patient associations, scientific reproductive societies, government institutions, National Health Service, European and international registries on ART. Moreover, there is a lot of information to better understand problems related to infertility reasons, news on reproductive and infertility issues, and a steady overview on Italian and European legislation on the reproductive field.
- A service for all centres: they can fill the forms on their activity each year and they have access to their local authority and to the national Register staff. They can also see on their reserved area how they perform in respect with some national indicators.
- A service for all 20 Italian Regions: They can see all the data relating the centres operating on their territory and they can monitor and elaborate data on their specific activity.

The Registry's website, last year, was visited by approximately 140.000 users, with a daily average of about 380 hits, and is the second most visited site in the National Institute of Health Portal.

1. ACCESS AND UTILIZATION OF ART SERVICES IN ITALY, 2018

1.1. Access to ART service

Figure 1 shows the regional distribution of the 202 ART authorized centers in 2018, but only 188 performed at least one ART cycle.

The largest number of ART centers is concentrated in Northern Italy (81 centers the 40,1% of the total) and in the Southern area (73 centers the 36,1% of the total), irrespective of the amount of their activity.




Figure 1. Regional distribution of the ART authorized centers and in brackets the number of centers per 100.000 women of reproductive age (15-45 years)*, 2018

*Average resident population in Italy in 2018: source ISTAT.

Table 1 shows the geographical distribution of ART centers according to the type of services offered. Overall, the number of centers active in 2018 was 202, 91 of which (45,0%) operating within the National Health Service (public and private), and 111 (55,0%) which provided only private service. The higher percentage of ART centers providing public service was concentrated in the North of Italy, i.e. in the North West 78%, while in the Centre and in the South of Italy there were mainly private facilities (52,5% and 72,6%, respectively).

Table 1. ART authorized centers distribution by region and type of service, 2018

Region and Geographical Area	ART centers	Art centers by type of Service					
		Public		Private covered by NHS		Private	
		N	%	N	%	N	%
Piemonte	13	5	38,5	1	7,7	7	53,8
Valle d'Aosta	1	1	100	0	0	0	0
Lombardia	25	13	52,0	10	40,0	2	8,0
Liguria	2	2	100	0	0	0	0,0
North-West	41	21	51,2	11	26,8	9	22,0
A.P. Bolzano	3	1	33,3	0	0	2	66,7
A.P. Trento	1	1	100	0	0	0	0
Veneto	18	8	44,4	0	0,0	10	55,6
Friuli Venezia Giulia	3	2	66,7	1	33,3	0	0
Emilia Romagna	15	6	40,0	0	0	9	60,0
North-East	40	18	45,0	1	2,5	21	52,5
Toscana	15	5	33,3	5	33,3	5	33,3
Umbria	2	1	50,0	0	0	1	50,0
Marche	3	2	66,7	0	0	1	33,3
Lazio	28	6	21,4	1	3,6	21	75,0
Central	48	14	29,2	6	12,5	28	58,3
Abruzzo	4	2	50,0	0	0	2	50,0
Molise	1	0	0	0	0	1	100
Campania	27	7	25,9	0	0	20	74,1
Puglia	11	2	18,2	0	0	9	81,8
Basilicata	1	1	100	0	0	0	0
Calabria	4	0	0,0	0	0	4	100
Sicilia	22	5	22,7	0	0	17	77,3
Sardegna	3	3	100	0	0	0	0
South and Islands	73	20	27,4	0	0	53	72,6
Italy	202	73	36,1	18	8,9	111	55,0

1.2. Utilization of ART services

Table 2 shows the time-trends of ART initiated cycles per million inhabitants and per million women of reproductive age (between 15 and 45 years), in comparison with the same indicators for Europe. Since 2005 In Italy, both the indicators were constantly growing, with an increase of 661 cycles (+103,9%) and of 4.658 cycles (+173,6%), respectively. The latest European data available refers to the activity of year 2016¹.

The number of started cycles per million inhabitants (calculated only for the 18 countries that have reported data of 100% of the centers) was 1.410 cycles in Europe versus 1.237 in Italy, both under the proposed optimal level of demand calculated as 1.500 of ART services per million inhabitants per year².

Table 2. Number of initiated ART cycle per million inhabitants and per million women of reproductive age (15-45 years) annually in Italy (2005-2018) and in Europe (2005-2016)

Years	ART Cycles/ million population		ART cycles/ million women (15 - 45 years)	
	Italy	Europe ^a	Italy	Europe ^a
2005	636	1.115	2.683 ^b	4.008 ^b
2006	692	850	3.328	3.503
2007	736	886	3.569	4.320
2008	800	947	3.905	4.661
2009	865	1.067	4.265	5.455
2010	973	1.221	4.863	6.258
2011	1.063	1.269	5.392	6.556
2012	1.078	1.252	5.562	6.519
2013	1.070	1.175	5.601	6.210
2014	1.102	1.399	5.855	7.608
2015	1.175	1.432	6.341	7.795
2016	1.237	1.410	6.781	7.794
2017	1.275	-	7.106	-
2018	1.297	-	7.341	-

a: data for Europe refers only to those country where data coverage was 100% in every year.

b: in 2005 ART cycles are related to the number of women aged between 15 and 49 years.

Figure 2 shows the distribution of initiated cycles per million women of childbearing age per geographical region. There is a great difference in the number of cycles performed among regions. If we select regions with more than five hundred thousand women in reproductive age, the distribution of cycles range from 3.642 cycles offered in Puglia region to 14.769 cycles provided in Toscana region. More in general, only 7 regions in Northern and Central area have numbers above the national average (7.341 cycles), while all the Southern regions have numbers below the average.




Figure 2. Regional distribution of the number of initiated ART cycles per million women of reproductive age (15-45 years)* and in brackets the percentage of initiated cycles performed in patients coming from other regions, 2018

*Average resident population in Italy in 2018: source ISTAT.

Since its establishment, IARTR collected data on 158.465 infants, of which 130.961 born from ART and 27.504 born from IUI cycles. These data have to be considered with caution due to the proportion of pregnancies lost to follow-up, that however changed to the better: from 21,5% in 2006 to 9,7% in 2018. In particular, in 2018, a number of 12.646 babies are born alive from ART techniques in Italy, that represent the 2,9% of the national babies born in the country in the same year. These values are an expression of the application of the ART techniques in the different regions, but they do not faithfully reflect the number of births because the cycles carried out also include the techniques performed on patients coming from outside the region (for example in Toscana region, 57,5% of cycles are performed on patients coming from other regions).




Figure 3. Regional distribution of the babies born in 2018 from ART cycles, also with donation, and the percentages in relation to the annual number of children born*

*Number of live babies born in Italy in 2018: source ISTAT.

Figure 4 shows the percentages of live-born babies conceived by ART compared with the national total number of children born in Italy. The percentage of infants born with ART procedures increased more than 4 times from 2005 to 2018.




Figure 4. Time-trends of the percentage of live babies born after ART also with donation cycles and after ART & IUI also with donation cycles in relation to the annual national number of children born in Italy, 2005-2018

2. EFFICACY OF ART IN 2018 AND TIME TRENDS FROM 2005 TO 2018

2.1. Overview of ART

2.1.1. What types of ART cycles were performed?

When an ART treatment is applied not using cryopreserved oocytes or embryos, is defined as a “**fresh cycle**”, and it includes:

- **In Vitro Fertilization with embryo transfer (IVF)**: an ART procedure that involves extracorporeal fertilization of gametes³;
- **Intra Cytoplasmic Sperm Injection (ICSI)**: a procedure in which a single spermatozoon is injected into the oocyte cytoplasm³.

An ART treatment in which cryopreserved oocytes or embryos are utilized is defined as a “**frozen/thawing cycle**”, and it includes:

- **Frozen/thawed Embryo Transfer (FET) cycle**: ART procedure in which cycle monitoring is carried out with the intention of transferring to a woman a frozen/thawed or vitrified/warmed embryo(s)/blastocyst(s)³;
- **Frozen/thawed Oocyte (FO) cycle**: ART procedure in which cycle monitoring is carried out with the intention of fertilizing thawed/warmed oocytes and performing embryo transfer³.

An ART treatment in which are used gametes that did not originate from the female recipient and/or her male partner is defined as a “**donation cycle**”, and it includes:

- **Oocyte donation (OD) cycle**: an ART cycle in which a woman receives oocytes from a donor to be used for reproductive purpose³;
- **sperm donation**: a cycle in which a woman receives spermatozoa from a person who is not her sexually intimate partner³;
- **double donation**: an ART cycle in which oocytes and spermatozoa both originating from donors are used.

The major part of the 79.734 Art cycle performed in Italy in 2018 were made with a fresh procedure (64,1%), then 24,6% of the cycles were performed with a FET, the 1,7% with a FO and the remaining 9,7% (7.743 cycles) with a gamete donation (see **Figure 5**).




Figure 5. ART cycles performed in 2018.

2.1.2. How did the types of ART techniques change according to transfers among women of different ages?

The patient's age is the variable that influences at the most the success of assisted reproduction techniques, and therefore the probability of obtaining a pregnancy. **Figure 6** shows the percentage of ART transfers performed in 2018 according to women's age. As expected, ART treatment with their own oocytes are the most applied techniques in women from ≤ 34 years till 40-42 years of age, while after 42 years of age the oocyte donation cycles reach 49,4% of application.




Figure 6. Types of ART procedure performed by female patients' age groups, 2018.

2.2. ART, non-donor cycles

2.2.1. What are the causes of infertility of couples in ART treatment using fresh cycles in 2018?

- *Female factor:*

- **tubal factor** - fallopian tubes are blocked or damaged, could prevent sperm from getting to the egg and eggs from getting to the uterus³;
- **ovulatory dysfunction** - ovaries are not producing eggs normally. The ovaries develop many small cysts instead of ripening and maturing one egg in each cycle.
- **endometriosis** – a disease characterized by the presence of endometrium-like epithelium and stroma outside the endometrium and myometrium. This condition can affect both fertilization of the egg and embryo implantation³.
- **diminished ovarian reserve** – indicate a reduced number and/or reduced quality of oocytes, such that the ability to reproduce is decreased³;
- **multiple abortions**-when there were two or more miscarriages without any full-term pregnancy.
- **multiple factor, female** - more than one female's cause of infertility.

- *Male factor* – abnormal semen parameters or function; anatomical, endocrine, genetic, functional or immunological abnormalities of the reproductive system; chronic illness; and sexual conditions incompatible with the ability to deposit semen in the vagina³.

- *Male and female factors* - one or more female and male causes of infertility.

- **Genetic factor** - Due to chromosomal abnormalities (numerical and/or structural) or to genetic alterations. They can be both male and female factors

- *Unexplained infertility* - no cause of infertility is found in either woman or man³.




Figure 7. Causes of infertility of couples in ART treatment using fresh cycle, 2018

2.2.2. What are the steps for an ART treatment using fresh cycles?

An ART cycle using fresh gametes:

- it is started when a woman begins taking fertility drugs to stimulate the ovaries to produce eggs or having her ovaries monitored for follicle production, if no drugs are given (**initiated cycle**)
- It continues, if the egg follicles are produced, by an ovarian follicular aspiration performed with the aim of retrieving oocytes (**retrieval**)
- After eggs collection, a process initiated by entry of a spermatozoa into a mature oocyte followed by formation of the pronuclei (**fertilization**)
- If fertilization occurs, the embryo(s) are cultured from day 1 to day 7, and then embryo(s) (generally 1 or 2) is(are) transferred to the woman's womb either at day 2 to 3 (cleavage stage) or at day 5 to 7 (blastocyst stage) (**transfer**)
- The attachment and subsequent penetration by a zona-free blastocyst into the endometrium (**implantation**)
- If implantation is successful, a pregnancy diagnosed by ultrasonographic visualization of one or more gestational sacs or definitive clinical sign of pregnancy occurs (**clinical pregnancy**)
- Generally in 75-80% of cases a **live birth delivery** occurs. A birth of twins, triplets or more are counted as one live birth.

Figure 8 shows outcomes resulting from various steps of fresh cycles performed in 2018. Of the 51.086 fresh cycles, 90,8% resulted in an egg retrieval, 59,9% in an embryo transfer, and 16,1% in freeze-all cycles. The “freeze-all” strategy could be decided for different reasons: either to avoid the risk of OHSS (Ovarian Hyper Stimulation Syndrome) or to perform PGT (Pre-implantation Genetic Testing). Because of the “freeze-all cycles”, we decided to compute the pregnancy rate per cycle removing them from the results of the fresh cycles. They will be included only when we computed the cumulative pregnancy rate (see Figure 11 and 20). Among the outcomes we must also consider the 10,9% of pregnancies lost to follow-up and the 26,3% of monitored pregnancies that do not reach the delivery due to miscarriages or ectopic pregnancies (see **Figure 12**).




Figure 8. Outcomes of ART using fresh cycles by stage of the treatment cycle, 2018

2.2.3. Did the number of embryos transferred differ among women of different age groups?

Figure 9 shows the distribution of the ART non donor transfers divided by the number of embryo(s) transferred according to women's age (due to the aggregate data collection we cannot have this information for each ART techniques). The number of transfers with three embryos increased with women's age when they use their own oocytes, while the number of transfers performed with one or two embryos declined with women's age. The single embryo transfer was performed in the 52.5% of transfers in women under 35 years and in the 43,5% in women older than 42 years.




Figure 9. Distribution of embryo transfers according to the number of embryos transferred (ART total) by female patients' age groups, 2018

2.2.4. What are the percentages of initiated cycles or thawings, and transfers that result in pregnancies for ART cycles?

Figure 10 shows the pregnancy rates per aspiration/thawing and per transfer for the ART techniques. Overall, the rates after FET (Frozen Embryo Transfer) were higher than others techniques, while the FO (Frozen Oocytes) ones were the lowest. The better results obtained with frozen embryos could be partially due to the selection of “good prognosis” patients (freeze-all strategy and/or more efficient single embryo transfer) in cryopreservation techniques. These results are observed also at the European level¹.




Figure 10. Pregnancy rates per aspiration/thawing and per transfer according to different ART procedures, 2018

2.2.5. What is the “Cumulative Pregnancy Rate”?

The cumulative pregnancy rate (CPR) is the overall chance of obtain a pregnancy from all fresh and frozen embryo transfers coming from one egg retrieval. Individual data instead of summary data collection would be needed to precisely calculate the CPR. Unfortunately, IARTR as well as other Registries around the world collects information only in aggregated form. To overcome these limitations and calculate CPR using aggregated data like EIM and ICMART, we sum the number of pregnancies obtained from fresh and frozen cycles divided by the number of aspirations, per year. CPR may provide a broader view of pregnancies that are achieved in Italy, in a year of activity. Moreover, the comparison of pregnancy rates from fresh cycles vs. cumulative pregnancy rates may show the estimated added value of embryo and oocyte cryopreservation.

Figure 11 shows pregnancy rate per fresh cycle and cumulative pregnancy rate by women age groups. Overall, embryo and oocyte cryopreservation increased by 75% the chances of achieving a pregnancy per started cycle. It should be emphasized that as we cannot take into account the weight of the "freeze-all" policy in every age group of female patients (data not collected by the Registry), so the pregnancy rates per initiated cycle could be underestimated in each age group.




Figure 11. Pregnancy rates per aspiration and cumulative pregnancy rates per aspiration, by female patients' age groups, 2018

2.2.6. What percentage of ART pregnancies resulted in a delivery?

Figure 12 shows the outcomes of the monitored clinical pregnancies obtained from the application of ART non donor procedures in Italy in 2018. Of the 7.401 monitored pregnancies issued from fresh cycles 73,7% resulted in a delivery, the 10,3% in multiple deliveries and the 24,6% in a miscarriage. In FET and FO procedures the 73,7 % and 73,6%, respectively, resulted in a delivery. The percentages of multiple deliveries in FET and FO are lower (5% and 9,1% respectively) then in fresh cycles, while the percentage of miscarriages were no significantly different.




Figure 12. Comparison of the outcomes of monitored clinical pregnancies that resulted from ART non donor procedures, 2018

2.2.7. What is the gender distribution of infertility factors among ART users?

Figure 13 shows major causes of infertility among patients who had ART using fresh cycles in 2018. Diagnoses range from one infertility factor in the patient or partner to multiple infertility factors in either one or both members of the couple. The male factors are decreasing since 2009, while the female ones are increasing since 2015. Overall, considering also the causes of infertility present in both members of the couples, female factors were diagnosed in more than 50% of couples in each year of data collection.




Figure 13. Time-trends of gender distributions of infertility causes among patients who had ART using fresh cycle, 2018. Total couples treated 42.090

2.2.8. Is the use of ART procedures change over time?

Figure 14 shows the number of ART non-donor initiated/thawing cycle and the number of ART active centers (with at least 1 cycle performed in the year) from 2005 to 2018.

The numbers of initiated cycles increased by 93% over time with an average increase of 7,2% for each year and it seems not to be related to the number of active centers.

Intra-Cytoplasmic Sperm Injection (ICSI) was originally developed to improve fertilization rates in couples with severe male factor infertility indication. Today, this procedure is widely used even without a reported diagnosis of male factor infertility.

The number of ICSI cycles increased from 24.209 in 2005 to 43.215 in 2018, while IVF cycles decreased from 8.994 to 7.871. In addition, FET thawing increased from 1.338 to 19.587, while FO procedures decreased from 2.711 to 1.318.




Figure 14. Time-trends of ART non donor initiated cycles/thawings and number of active ART centers, 2005–2018

2.2.9. Did the use of FET (Frozen Embryo Transfer) procedures differ in Italy compared to the other European countries over time?

In 2004, the Italian Parliament approved a Law (40/2004) regulating ART in which embryo cryopreservation was banned. In 2009 Italian Constitutional Court removed some limitations set out in the Law, including the practice of embryo freezing, now permitted under specific conditions. For this reason the use of FET has declined consistently after 2004 and resumed steadily after 2009 (**Figure 14**). As shown in **Figure 15** percentage of FET cycles performed highly increased from 3,6% in 2005 to 27.2% in 2018. In comparison with some European countries¹, such as France, Germany and Sweden, for 2016 last European data available, Italy still shows the lowest number of FET cycles performed, but they are steadily increasing.




Figure 15. Time- trends of percentages of FER thawing on total ART non donor initiated cycles in selected European countries (2005-2015)* and Italy (2005-2018)

*Elaboration of number of ART treatment cycles: source ESHRE reports on ART.

2.2.10. Has the age of ART female patients changed over time?

Figure 16 shows the distribution of fresh cycles by women age groups, from 2005 to 2018.

For women older than 40 the percentage of fresh cycles performed increased from 20,7% in 2005 to 34,6% in 2018, whilst the percentage of fresh cycles performed in women ≤ 34 years old decreased from 39,3% in 2005 to 27,6% in 2018. Overall, the mean age of women who had fresh cycles increase over time from 35,3 to 36,7 years.




Figure 16. Time-trends of fresh initiated cycles distributions by age classes of female patients, 2005-2018

2.2.11. Has the number of embryos transferred changed in fresh cycles?

Figure 17 shows trends with the percentage of the number of embryos transferred in fresh cycles. From 2005 to 2018 the single and the double embryo transfer increased from 18,7% to 39,1% and from 30,9% to 49,4%, respectively. On the other hand, transfers with 3 embryos dramatically decreased from 50,4%, first to 38,1% in 2010 to reach 10,9% in 2018. As shown in the figure, this trend begins from the end of 2009 when Law 40/2004 was changed, and the limit to transfer all the embryos created for a maximum of three removed. Values of transfers with four or more embryos has decreased to 0,6% in 2018, after having been quite stable during time. The average number of embryos transferred decreased from 2,3 embryos per transfer in 2005 to 1,8 in 2018. This trend is similar in most European countries¹.




Figure 17. Time-trends of transfer by number of embryos transferred, 2005-2018

2.2.12. Did pregnancy rates per transfer changed over time among different ART procedures?

Figure 18 shows pregnancy rates per transfer in order to compare cycles with fresh oocytes and embryos vs. those using frozen embryos (FET) or frozen oocytes (FO).

Overall, FET cycles showed the best pregnancy rates increasing highly from 16,3% in 2005 to 31,5% in 2018, those with fresh oocytes slightly increased from 24,5% to 27,2%, and those with frozen oocytes from 11,4% to 21,5%. Since five years, FET pregnancy rates are higher than the fresh ones.




Figure 18. Time-trends of pregnancy rate per transfer for fresh, thawed embryos (FET) and thawed/warmed oocytes cycles (FO), 2005-2018

2.2.13. Did cumulative pregnancy rates per initiated cycle changed over time?

Figure 19 compares the percentage of pregnancies obtained on fresh initiated cycles with the Cumulative Pregnancy Rate (CPR) over time. As described in **Chapter 2.2.5**, CPR is presented as the sum of pregnancies obtained from fresh cycles and with frozen/thawing cycles (FET and FO) as nominator and the number of initiated cycles with fresh techniques of the same year as denominator. CPR gives an estimate of the likelihood of obtaining a pregnancy for a woman undergoing an ART cycle, also having the opportunity to perform oocyte and/or embryo thawing cycles. Moreover, the comparison of pregnancy rate from fresh cycles vs. CPR may show the estimated benefit of embryo and oocyte cryopreservation. As the figure shows, this value is constantly growing from +6,6% in 2005 to a +49,5% in 2018.




Figure 19. Time-trends of pregnancy rate per initiated cycle for fresh and cumulative pregnancy rate per initiated cycle, 2005-2018

2.2.14. Does the risk of pregnancy loss differ among women of different age groups?

Increasing female age also increases the risk of negative pregnancy outcomes (spontaneous or therapeutic abortions and ectopic pregnancies). As shown in **figure 20** pregnancy loss rates were much higher in older age groups. Rates decreased over time from 65,2% to 57,8% for women older than 43 and from 42,5% to 40,9% for women aged 40-42. Rates for women younger than 40 were quite stable over time.




Figure 20. Time-trends of percentages of total pregnancy loss using ART non donor cycles by female age groups, 2009-2018

2.3. ART donor cycles

In April 2014 Italian Constitutional Court removed the prohibition (set out in the Law 40/2004), regarding the practice of ART techniques using donor gametes. Currently oocyte donation, semen donation and double donation are allowed.

See the Summary table ([see page 55](#)) for more detailed data on activity and outcomes regarding ART donor cycles in 2018.

2.3.1. Which gametes were used in ART donor cycles in 2018?

Figure 21 shows the distribution of the types of the 7.743 ART cycles using donor gametes applied in Italy in 2018 and the origin of the donated frozen oocytes. In about 82,7% of ART donor cycles, fresh or frozen eggs were used, in 17,3% of cycles there is a donation of male gametes only. In the oocyte donation cycles, almost all cycles (98,2%) were performed with oocytes obtained from a foreign bank.




Figure 21. Distribution of all ART cycles using donor gametes and origin of the oocytes for the donor oocyte cycles, 2018. Total cycles = 7.743

2.3.2. What was the age of recipient female patients in ART donor cycles in 2018

Figure 22 shows the distribution of performed transfers according to the recipient female age group at the start of a donation cycle. The different age distribution in the different types of gametes utilised reflects the treatment indication.




Figure 22. Distribution of transfers cycles using donor gametes by recipient female age groups, 2018

2.3.3. What was the percentage of transfers that result in pregnancies in ART donor cycles in 2018?

Figure 23 shows the pregnancy rate per transfer in ART donor techniques.




Figure 23. Distribution of pregnancy rate per transfer using donor gametes, 2018

2.3.3. What were the outcomes of pregnancies obtained in ART donor cycles in 2018?

Figure 24 shows the outcomes of the 2.171 monitored clinical pregnancies obtained from the application of ART donor procedures in Italy in 2018. Of the 357 monitored pregnancies obtained from sperm donation 10,6% resulted in multiple deliveries and 22,1% in an abortion. In frozen donor oocyte cycles the multiple deliveries rate was higher (11,5%) than in fresh donor oocyte ones (5,4%). While there were no differences between the types of donation for the percentage of abortion: 22,1% for sperm donation cycles, 20,6% for fresh oocytes, 22,1% for frozen oocytes donation cycles and 21,1% for double donation cycles.




Figure 24. Comparison of the outcomes of monitored clinical pregnancies that resulted from ART donor cycles, 2018

2.4. PGT (Pre-Implantation Genetic Testing) activity

During an ART treatment, some investigation could be performed to analyze the DNA from embryos (at cleavage stage or blastocyst) for HLA-typing or for determining genetic abnormalities. These includes:

- **PGT-A** - test for the detection of aneuploidies;
- **PGT-M** – test for monogenic/single defects;
- **PGT-SR** – tests for chromosomal structural rearrangements.

2.4.1. Which types of genetic tests were performed in ART cycles in 2018?

PGT activity, recorded from 46 centers, involved 2.447 tests (2.352 fresh and 95 thawing). They result in 1.825 fresh and 25 frozen embryo transfers. **Figure 25** shows the distribution of the application of PGT (Preimplantation Genetic Testing) cycles in Italy. A total of 825 pregnancies (45% per transfer) and 688 deliveries (36,9% per transfer) resulted from fresh cycles, while 10 pregnancies (40% per transfer) and 6 deliveries (24% per transfer) resulted from frozen embryo cycles. Please check on summary table ([see page 54](#)) for more detailed data on activity and outcomes regarding PGD/PGS cycles in 2018.




Figure 25. Distribution of PGD/PGS tests, 2018

2.4.2. Did the use of different genetic analysis in ART change over time?

Since the first data collection on PGT cycles performed on 2014 activity, the number of centers performing at least one cycle increased from 22 in 2014 to 46 in 2018. At the same time also the number of analysis performed increased from 1.695 cycles in 2014 to 2.447 cycles in 2018 (Figure 26).




Figure 26. Time-trend of number of PGT tests and of n° of ART centers performing at least one PGD/PGS tests, 2018

3. ART SAFETY INDICATORS

3.1. Safety in ART procedures

Although ART techniques are considered a safe medical procedure, in few cases some complications could arise during the treatment. These complications could arise at the moment of the ovarian stimulation or during pick-up procedure. Both could affect women's health. Another kind of complication for mother and child it is strictly related to embryo transfer policies. The transfer of more than one embryo could determine multiple pregnancies, therefore multiple deliveries and multiple births that could determine prematurity, morbidity and perinatal mortality of the babies.

3.1.1. Did the numbers of complications for ART cycles change over time?

Complications in an ART treatment are considered:

- **Ovarian Hyper Stimulation Syndrome (OHSS):** An exaggerated systemic response to ovarian stimulation characterized by a wide spectrum of clinical and laboratory manifestations. It may be classified as mild, moderate or severe according to the degree of complications³. It is registered as a complication in the ART Register when it is diagnosed as "severe" (at least grade 3).
- **Bleeding:** Significant bleeding, internal or external, after oocyte aspiration retrieval requiring hospitalization for blood transfusion, surgical intervention, clinical observation or other medical procedure³.
- **Infection:** The presence of a bacterial or viral infection that can occur during any surgical procedure.

Cases of bleedings and infections are decreasing during time, while the number of OHSS is quite stable.




Figure 27. Time-trends of complications in fresh non donor ART treatments, 2005-2018.

3.1.2. Did the percentages of multiple deliveries for ART non donor cycles change over time?

Figure 28 shows trends for multiple deliveries in fresh cycles.

From 2005 to 2018 twin delivery percentage decreased from 21,6% to 13,6% while percentages of triplets and more deliveries decreased from 2,7% to 0,3%. This value is similar to the average European value of 0,3% reported in 2016 EIM data¹. We should remember that from 2004 till 2009 the Law obliged to transfer at once, all the embryos created for a maximum of three.




Figure 28. Time-trends of deliveries from fresh cycles according to gestational order, 2005-2018.




Figure 29. Time-trends of deliveries from FER thawing according to gestational order, 2005-2018




Figure 30. Time-trends of deliveries from FO thawing according to gestational order, 2005-2018

3.1.2. Did the percentages of preterm live babies change over time?

Figure 31 shows trends of ART preterm live born babies that are highly correlated with the multiplicity of deliveries.

The percentage of preterm live babies in singleton and twin deliveries are quite stable during all the period. Otherwise, in triplet deliveries, there is a variability from year to year, but overall the trend is upwards from 82,7% in 2005 to 96,1% in 2018 while it is stable or slightly lower in singletons.




Figure 31. Time-trends of percentage of preterm ART live born babies (<37 week of gestation) by gestational order, 2005-2018

3.1.4. Did the percentage of underweight live babies change over time?

Figure 32 shows the trends of ART live born babies underweight that are highly correlated with the multiplicity of deliveries as already described in respect to prematurity.

In babies born underweight, the percentage in singleton and twin deliveries are quite stable during all the period. Overall, the trend is also quite stable for triplet and more deliveries from 90% in 2005 to 88,5% in 2018 with a minimum value of 63.3% in 2014 due to variability of the small number.




Figure 32. Time-trends of percentage of ART live born babies underweight (<2,500 gr) by gestational order, 2005-2018

4. INTRA-UTERINE INSEMINATION PROCEDURES

4.1. Access to Intra-Uterine Insemination service

Figure 33 shows the regional distribution of the 345 Intra-Uterine Insemination (IUI) authorized centers in 2018, but only 291 performed at least one homologous IUI cycle.

The majority of centers is concentrated in Northern Italy (159, 46,1% of the total) and then in the South (119 centers, 34,5% of the total), irrespective of the amount of activity they performed . Even with some differences the access is almost good in all the regions.




Figure 33. Regional distribution of IUI + ART authorized centers and in brackets the number of centers per 100000 women of reproductive age (15-45 years)*, 2018

*Average resident population in Italy in 2018: Source ISTAT.

As shown in **Table 3**, in 2018 there were 345 authorized centers that perform IUI of which only 126 operating within the National Health System (public and private 36,5%) and 219 providing private service (63,5%). Most of public centers that could perform IUI in Italy were in the North: 62 out of 106 centers (58,5%).

Table 3. IUI authorized centers distribution by region and type of service, 2018

Region and Geographical Area	Total	Type of Service					
		Public		Private covered by NHS		Private	
		N	%	N	%	N	%
Piemonte	23	8	34,8	1	4,3	14	60,9
Valle d'Aosta	1	1	100	0	0,0	0	0,0
Lombardia	58	18	31,0	10	17,2	30	51,7
Liguria	9	5	55,6	0	0,0	4	44,4
Northwest	91	32	35,2	11	12,1	48	52,7
P.A. Bolzano	6	4	66,7	0	0,0	2	33,3
P.A. Trento	1	1	100	0	0,0	0	0,0
Veneto	35	13	37,1	2	5,7	20	57,1
Friuli Venezia Giulia	4	3	75,0	1	25,0	0	0,0
Emilia Romagna	22	10	45,5	0	0,0	12	54,5
Northeast	68	31	45,6	3	4,4	34	50,0
Toscana	23	8	34,8	5	21,7	10	43,5
Umbria	2	1	50,0	0	0,0	1	50,0
Marche	6	2	33,3	0	0,0	4	66,7
Lazio	36	6	16,7	1	2,8	29	80,6
Central	67	17	25,4	6	9,0	44	65,7
Abruzzo	6	3	50,0	0	0,0	3	50,0
Molise	2	0	0,0	0	0,0	2	100
Campania	44	9	20,5	0	0,0	35	79,5
Puglia	15	3	20,0	0	0,0	12	80,0
Basilicata	2	2	100	0	0,0	0	0,0
Calabria	9	1	11,1	0	0,0	8	88,9
Sicilia	38	5	13,2	0	0,0	33	86,8
Sardegna	3	3	100	0	0,0	0	0,0
South and Islands	119	26	21,8	0	0,0	93	78,2
Italy	345	106	30,7	20	5,8	219	63,5

4.2. Efficacy safety and trends of IUI

4.2.1. Is the use of IUI-H increasing?

Intra-Uterine Insemination (IUI) is a medical procedure in which a laboratory processed sperm is placed into the women’s uterus to attempt a pregnancy³. It can be performed using husband semen (**IUI-Homologous**) or with the semen of an anonymous donor (**IUI-Donor**).

As described for ART techniques using donor gametes ([see page 30](#)) the IUI-D procedures have been permitted only after the Italian Constitutional Court sentence in April 2014. For the activity of IUI-D in 2018, please check on Summary table for 2018 ([see page 55](#)) for more detailed data.

Figure 34 shows the use of IUI-H from 2005 to 2018. The number of IUI-H cycles decreased from 26.292 to 17.083 after having reached his maximum of 33.335 cycles in 2009. There were no changes in pregnancy rate from 10,7% in 2005 to 10,6% in 2018. Average women age increases of 0,4 year during time.




Figure 34. Time-trends of outcomes of IUI-H cycles, 2005-2018

4.2.2. Do percentages of IUI-H cycles resulting in pregnancies differ among women of different age groups?

Figure 35 shows percentages of insemination cycles for IUI-H that resulted in pregnancies and in deliveries among women of different age groups.

The probability to obtain a pregnancy and to reach a delivery in an IUI-H treatment is highly related to the age of women. The percentage in older women are very low: 3,3% for pregnancy and 1,9% for delivery in over 43.




Figure 35. Pregnancy rates and delivery rates per insemination for IUI-H cycles by age groups of female patients, 2018

4.2.3. What were the outcomes of pregnancies obtained in IUI-H cycles in 2018?

Figure 36 shows the outcomes of the monitored clinical pregnancies obtained from the application of the homologous IUI in Italy in 2018. Of the 1.607 monitored pregnancies in homoulogous IUI cycles 7,0% resulted in multiple deliveries and the 19,2% have a miscarriage.




Figure 36. Outcomes of clinical pregnancies that resulted from H-IUI cycles, 2018

4.2.4. Did the numbers of complications for H-IUI cycles change over time?




Figure 37. Time-trends of complications in homologous-IUI treatments, 2005-2018

4.2.5. Did the percentages of multiple deliveries change over time for homologous intrauterine insemination cycles?

Figure 38 shows time trends for multiplicity of deliveries in IUI-H cycles.

From 2005 to 2018 twin deliveries rates decreased from 15% to 8,4% while percentage of triplets and more deliveries are quite stable. Since 2007, 9 deliveries out of 10 were singleton.




Figure 38. Time-trends of deliveries from IUI-H cycles according to gestational order, 2005-2018

4.3. IUI donor cycles.

4.3.1. What was the outcome in IUI donor cycles in 2018?

In total 691 initiated cycles using donor sperm in IUI cycle started in 2018, 20,1% resulted in a clinical pregnancy and 13,5% resulted in a delivery. However, most of these cycles (about 80%) did not produce a pregnancy while a small proportion (3,3%) resulted in a pregnancy loss (i.e. ectopic pregnancy or miscarriage).




Figure 39. Outcome of IUI-D cycles, 2018 (691 initiated cycles)

APPENDIX
SUMMARY TABLE OF ACTIVITY AND
OUTCOME OF ART PROCEDURES
YEARS 2011-2018

Summary table of activity and outcome of all ART procedures, 2011-2018

	2011	2012	2013	2014	2015	2016	2017	2018
N° Clinics	354	355	369	362	366	360	366	345
% of clinics reporting data to ISS	100	100	100	100	100	100	100	100
ALL PROCEDURES (IUI-H, IUI-D, ART Non donor and Donor)								
N° Patients	73.570	72.543	71.741	70.826	74.292	77.522	78.366	77.509
N° Initiated cycles	96.427	93.634	91.556	90.957	95.110	97.656	97.888	97.149
N° Live born	11.933	11.974	12.187	12.720	12.836	13.582	13.973	14.139
ART activity (Fresh non donor, Thawing non donor, donor)								
N° Patients	53.558	54.458	54.523	55.859	59.747	63.724	65.943	66.083
N° Initiated cycles	63.783	64.207	64.447	67.054	72.048	75.889	78.457	79.735
N° Pregnancies	12.221	12.646	12.775	13.642	14.391	15.405	16.793	17.402
Cumulative Pregnancy Rate per fresh initiated cycle	21,8	22,8	23,2	24,5	26,0	28,6	31,7	32,8
% Pregnancies lost to follow-up	12,2	13,2	10,3	10,8	11,3	10,2	11,9	9,7
N° Deliveries	8.003	8.127	8.495	9.252	9.512	10.386	11.094	11.428
N° Live born	9.658	9.818	10.217	11.037	11.029	11.791	12.454	12.646
ART SERVICES AVAILABILITY INDICATORS								
ART Initiated cycles per 1 million women aged 15 and 45	5.293	5.562	5.601	5.860	6.341	6.781	7.106	7.341
ART Initiated cycles per 1 million inhabitants	1.050	1.078	1.070	1.103	1.175	1.237	1.275	1.297
IUI-H and IUI-D activity								
N° Patients	20.012	18.085	17.218	14.967	14.545	13.798	12.423	11.426
N° Initiated cycles	32.644	29.427	27.109	23.903	23.062	21.767	19.431	17.774
N° Pregnancies	3.246	3.024	2.775	2.399	2.466	2.429	2.078	1.952
% Pregnancy Rate per cycle	9,9	10,3	10,2	10,0	10,7	11,2	10,7	11,0
% Pregnancies lost to follow-up	18,1	17,1	16,8	18,2	16,8	15,0	13,9	11,7
N° Deliveries	2.062	1.974	1.810	1.530	1.649	1.629	1.396	1.369
N° Live born	2.275	2.156	1.970	1.683	1.807	1.791	1.519	1.493

Summary table of activity and outcome of ART non donor procedures, 2011-2018

	2011	2012	2013	2014	2015	2016	2017	2018
FRESH CYCLES (Non Donor)								
N° Patients	46.491	46.491	46.433	45.985	45.689	44.965	44.279	42.090
Average age calculated	36,48	36,50	36,55	36,68	36,68	36,80	36,70	36,74
N° Initiated cycles	56.092	55.505	55.050	55.705	55.329	53.906	53.014	51.087
N° aspirations	50.290	50.096	50.174	50.794	50.214	48.756	47.911	46.387
N° transfers	42.331	41.822	40.696	39.768	37.975	36.038	33.832	30.584
N° Pregnancies	10.959	11.077	10.712	10.834	10.081	9.326	9.310	8.307
% Pregnancies rate per cycles	19,5	20,0	19,5	19,4	18,2	17,3	17,6	16,3
% Pregnancies rate per cycles without freeze-all strategy	19,9	20,2	20,2	20,7	19,9	19,1	20,0	19,0
% Pregnancies rate per aspirations	21,6	22,1	21,3	21,3	20,1	19,1	19,4	17,9
% Pregnancies rate per transfers	25,9	26,5	26,3	27,2	26,5	25,9	27,5	27,2
% Twin Pregnancies	18,8	18,9	19,4	19,5	17,0	15,6	15,0	13,1
% Triplet or more Pregnancies	1,8	1,8	1,6	1,4	0,9	0,6	0,5	0,4
% Pregnancies lost to follow-up	12,7	13,9	10,9	11,9	13,1	11,2	13,4	10,9
N° Deliveries	7.193	7.116	7.125	7.277	6.498	6.196	6.029	5.458
N° Live born	8.734	8.680	8.677	8.848	7.695	7.172	6.951	6.186
FROZEN/THAWED EMBRYOS (FER)								
N° Patients	4.712	5.937	6.316	8.139	10.557	12.485	14.441	16.067
Average age calculated	35,0	35,2	35,2	35,4	35,2	35,3	35,3	35,2
N° thawing cycles	5.184	6.513	7.428	9.501	12.903	14.990	17.281	19.587
N° transfers	4.808	5.496	6.818	8.851	11.849	14.328	16.673	19.016
N° Pregnancies	910	1.231	1.763	2.448	3.379	4.128	5.059	5.995
% Pregnancies rate per thawings	17,6	18,9	23,7	25,8	26,2	27,5	29,3	30,6
% Pregnancies rate per transfers	18,9	22,4	25,9	27,7	28,5	28,8	30,3	31,5
% Twin Pregnancies	14,4	12,7	10,6	11,2	9,8	9,1	7,7	7,0
% Triplet or more Pregnancies	0,5	0,6	0,7	0,4	0,4	0,3	0,1	0,1
% Pregnancies lost to follow-up	6,7	8,4	7,0	5,5	5,6	6,7	7,9	6,6
N° Deliveries	584	790	1.169	1.747	2.403	2.890	3.486	4.127
N° Live born	666	888	1.308	1.929	2.609	3.104	3.703	4.393
FROZEN/THAWED OOCYTES (FO)								
N° Patients	2.355	2.030	1.774	1.530	1.418	1.341	1.281	1.201
Average age calculated	34,7	35,0	35,0	35,1	35,2	35,1	35,2	35,4
N° thawing cycles	2.507	2.189	1.969	1.639	1.529	1.460	1.391	1.318
N° transfers	2.012	1.736	1.491	1.295	1.221	1.187	1.146	1.036
N° Pregnancies	352	338	300	273	254	238	235	223
% Pregnancies rate per thawings	14,0	15,4	15,2	16,7	16,6	16,3	16,9	16,9
% Pregnancies rate per transfers	17,5	19,5	20,1	21,1	20,8	20,1	20,5	21,5
% Twin Pregnancies	12,2	13,3	13,0	12,5	10,6	12,2	12,8	9,9
% Triplet or more Pregnancies	1,1	0,3	1,7	0,8	0,4	1,3	0,4	0,0
% Pregnancies lost to follow-up	10,5	6,8	6,0	7,0	5,9	4,6	4,7	6,7
N° Deliveries	226	221	201	179	170	154	165	153
N° Live born	258	250	232	199	193	177	190	172

Summary table of activity and outcome of ART with PGT analysis, 2014-2018

	2014	2015	2016	2017	2018
PGT activity					
N° clinics reporting data (with at least 1 patient treated)	22	33	35	42	46
N° Patients	1.596	1.799	2.247	2.459	2.653
N° Tests performed	1.695	1.894	2.242	2.483	2.447
Patient average age	36,1	36,2	36,0	36,0	36,2
N° Pregnancies	459	617	721	859	850
% Pregnancies rate per transfer	45,5	45,1	44,4	43,8	45,0
% Pregnancies lost to follow-up	1,5	1,5	2,5	3,3	1,9
N° Deliveries	383	514	582	693	694
N° Live born	398	529	599	705	705

Summary table of activity and outcome of ART procedures with gamete donation, 2014-2018

	2014	2015	2016	2017	2018
ART-Donor activity					
N° clinics reporting data (with at least 1 patient treated)	17	69	83	91	101
N° Patients	205	2.083	4.933	5.941	6.725
N° Initiated cycles	209	2.287	5.533	6.771	7.743
N° Pregnancies	87	677	1.713	2.189	2.517
% Pregnancies lost to follow-up	26,4	15,7	13,9	15,5	13,7
N° Deliveries	49	441	1.146	1.414	1.690
N° Live born	61	532	1.338	1.610	1.895
IUI-D activity (sperm donation)					
N° clinics reporting data (with at least 1 patient treated)	13	52	65	62	73
N° Patients	32	379	517	487	488
N° Initiated cycles	37	513	714	743	691
N° Pregnancies	7	103	137	154	139
% Pregnancies rate per cycles	18,9	20,1	19,2	20,7	20,1
% Pregnancies lost to follow-up	85,7	30,1	16,8	16,9	16,5
N° Deliveries	1	61	98	110	93
N° Live born	1	69	119	127	107

BIBLIOGRAPHY

- 1) Wyns, C. (2020, July 31). ART in Europe, 2016: results generated from European registries by ESHRE. *Human Reproduction Open*, 3, 1-17. doi:10.1093/hropen/hoaa032
- 2) ESHRE Capri Workshop Group (2001). "Social determinants of human reproduction." *Human Reproduction* 16(7), 1518-1526.
- 3) F. Zegers-Hochschild et al. (2017). "The International Glossary on Infertility and Fertility Care, 2017", *Human reproduction* 32 (9), 1786-1801. doi:10.1093/humrep/dex234.