

ENGLISH VERSION



Istituto Superiore di Sanità

Rapporto ISS COVID-19 • n. 24/2020

Interim guidance for the appropriate support of adrenal insufficiency in children during the current SARS-CoV-2 pandemic emergency

ISS COVID-19 Rare Diseases Working Group

Version of May 10, 2020

Interim guidance or the appropriate support of adrenal insufficiency in children during the current SARS-CoV-2 pandemic emergency

Version May 10, 2020

ISS COVID-19 Rare Diseases Working Group

Domenica Taruscio, Istituto Superiore di Sanità, Roma
Luigi Bertinato, Istituto Superiore di Sanità, Roma
Anna Ladogana, Istituto Superiore di Sanità, Roma
Francesco Gabbrielli, Istituto Superiore di Sanità, Roma
Maurizio Pocchiari, Istituto Superiore di Sanità, Roma
Paolo Salerno, Istituto Superiore di Sanità, Roma
Marco Silano, Istituto Superiore di Sanità, Roma

in collaboration with:

ISS COVID-19 Drugs Working Group

Monica Boirivant, Istituto Superiore di Sanità, Roma

and in collaboration with:

Alessandra Cassio

Programma di Malattie Endocrino-Metaboliche, Unità di Pediatria, Dipartimento di Scienze Mediche e Chirurgiche, Università di Bologna, Coordinatore italiano 2020-2021 (Pediatria) European Reference Network on Rare Endocrine Conditions (ENDO-ERN)
in rappresentanza della Società Italiana di Endocrinologia e Diabetologia Pediatrica (SIEDP)

Nella Augusta Greggio

Poliambulatorio Morgagni-Affidea, Padova. Già coordinatore pediatrico italiano (2017-2019) per European Reference Network on Rare Endocrine Diseases (ENDO-ERN) in rappresentanza della Società Italiana di Endocrinologia e Diabetologia Pediatrica (SIEDP)

English version edited by Nella Augusta Greggio and Laura Lee Cellai

Istituto Superiore di Sanità

Interim guidance for the appropriate support of adrenal insufficiency in children in the current SARS-CoV-2 infection emergency scenario. Version of May 10, 2020.

ISS COVID-19 Rare Diseases Working Group

2020, 9 p. Rapporto ISS COVID-19 n. 24/2020 – English version

This report provides ad interim guidance for the appropriate support of children and adolescents with adrenal insufficiency during the current SARS-CoV-2 pandemic emergency. Adrenal insufficiency is a potentially lethal disease, characterized by a deficit in the production or action of glucocorticoids, sometimes associated with insufficiency of mineralocorticoids and androgens. The main clinical symptoms of adrenal insufficiency include weakness, fatigue, anorexia, nausea and vomiting, abdominal pain, weight loss, postural hypotension and in some cases desire to eat salt. In general, the diagnosis and treatment of patients with adrenal insufficiency is very demanding. In particular, the presence of fever $\geq 38^{\circ}\text{C}$, acute infection, stress such as surgery and other acute conditions can alter the metabolic status to such an extent that, if not treated properly, adrenal crisis in many forms of adrenal insufficiency may occur. Adrenal crisis is a critical emergency requiring immediate action, appropriate treatment and hospitalization, because it is potentially lethal. In case of suspected or confirmed COVID-19, cortisone replacement therapy must be continued and, depending on clinical severity, increased appropriately to guarantee adequate hormonal and cardiovascular support during conditions of physiological stress. Contact with reference centres to ensure advice from specialists is highly recommended, also through the use of telemedicine systems.

The original Italian version of ISS COVID-19 Reports are available at: <https://www.iss.it/rapporti-covid-19>

The reports translated in English are available from: <https://www.iss.it/rapporti-iss-covid-19-in-english>

For information: domenica.taruscio@iss.it

Cite this document as follows:

ISS COVID-19 Rare Diseases Working Group. *Interim guidance for the appropriate support of adrenal insufficiency in children in the current SARS-CoV-2 infection emergency scenario. Version of May 10, 2020.* Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 24/2020 - English version).

The responsibility for scientific and technical data lies with the authors, who declare that they do not have any conflict of interest.

Editing and graphics: ISS Scientific Communication Unit (Sandra Salinetti and Paola De Castro)

© Istituto Superiore di Sanità 2020
viale Regina Elena, 299 – 00161 Roma



Table of contents

SARS-CoV-2 infection in children	1
Adrenal insufficiency and SARS-CoV-2	3
Introduction.....	3
Main causes of adrenal insufficiency.....	3
Operational guidance.....	6
a. Procedures for delivery of therapies	6
b. Procedures for carrying out clinical tests	6
c. Management of therapy during infection.....	6
d. Assessment of infectious risk	6
e. Preventive measures.....	6
f. Support of telemedicine systems.....	7
References	8
Useful links	9

SARS-CoV-2 infection in children

Compared to the level of knowledge on the adult population, the prevalence of SARS-CoV-2 infection in the paediatric population is unknown. Current available scientific evidence indicates that in children, COVID-19 is a much less dramatic illness with significantly lower mortality than in adults.

A first review of the data of children with COVID-19 was performed in China and concludes that of 2143 children with confirmed infection, only 112 (5.6%) developed a severe form of the disease (defined as the presence of hypoxia / dyspnoea) and 13 (0.6%) developed Acute Respiratory Distress Syndrome (ARDS) (1).

A review of the literature made in April 2020 confirmed these statistics, even though the accentuation of the bronchial texture and “frosted glass” opacity on radiological examination were noted even in asymptomatic children (2).

In April 2020, the CDC (Centers for Disease Control and prevention in USA) published a second study. Out of 149,760 cases of SARS-CoV-2 with positive nasopharyngeal swab PCR tests who were identified between 12 February and 2 April, only 2,500 were children/adolescents under the age of 18. The presence of clinical symptoms suggestive of COVID-19 (fever, cough and dyspnoea) was observed in 73% of the 2500 paediatric SARS-CoV-2 positives whereas 93% of the positive adults were symptomatic. The same document highlights a hospitalization rate of at least 5.7%, and an ICU hospitalization ranging between 0.58% and 2%. In the paediatric group, 62% of hospitalized children were aged 12 months or less and five of 15 infants needed intensive care. About 77% of hospitalized patients, including all ICU patients, had one or more concomitant condition, whereas only 12% of non-hospitalized children had concomitant conditions (3).

In a series of 41 Spanish paediatric patients with confirmed SARS-CoV-2 infection, 25 required hospitalization, 4 patients required intensive care and another 4 patients needed respiratory assistance via nasal cannulas (4).

In Italy, ISS data report that up until May 7, 2020, of the 25,452 deaths with positivity for SARS-CoV-2 infection, only three cases were in the 0-19 years age group (5).

In a series of 100 children with SARS-CoV-2 positive swabs who entered the emergency room of 17 Italian hospitals, only 52% of patients with fever had the other two symptoms suggestive of COVID-19 (cough and dyspnoea). Thirty-eight of the children required hospitalization, 9 of whom needed respiratory support (6 with pre-existing diseases). All 100 children in the series have recovered (6).

These data indicate an overall infection and mortality rate in children that is significantly below the 4% rate reported in adults with COVID-19, thus reducing the concerns of doctors and parents. However, more attention should be paid to children with less than one year of age and to those with concomitant conditions who show the symptoms of the infection.

Studies performed on pregnant mothers infected with SARS-CoV-2 exploring the relationship between maternal immunity and protection of the infant from infection, did not reach conclusive results (7). “Suspicious” symptoms such as fever, no response to therapy, difficulty breathing, cough and sleepiness in infants born from a SARS-CoV-2 positive mother should alert the parents and the paediatrician.

Patients with hypoadrenalism are more at risk of contracting infections (8-10) and, in case of suspected or confirmed COVID-19, cortisone replacement therapy must be continued, or, depending on clinical severity, increased appropriately to guarantee adequate hormonal and cardiovascular support during

conditions of physiological stress. Contact with reference centres to ensure advices from specialists is highly recommended, also through the use of telemedicine systems.

Adrenal insufficiency and SARS-CoV-2

Introduction

Adrenal Insufficiency (or hyposurrealism) is a potentially lethal disease, characterized by a deficit in the production or action of glucocorticoids, with or without deficiency in mineralcorticoids and adrenal androgens.

It can result from primary adrenal failure or secondary adrenal disease due to impairment of the hypothalamic-pituitary axis (central adrenal insufficiency). In the context of central adrenal insufficiency, we distinguish a secondary form caused by an insufficient pituitary ACTH production and a tertiary form due to an altered hypothalamic function with inadequate secretion of CRF (11).

The estimated incidence of the primary form is 4.4-6 new cases per million inhabitants per year. Tuberculosis was the most common cause of primary adrenal insufficiency in the first half of the 20th century, but today the most frequent form is that of autoimmune disease. The frequency of the various forms of primary insufficiency in children differs substantially from that of the adult population, with genetic forms being the most common in children, in particular the classic form of congenital adrenal hyperplasia due to 21-hydroxylase deficiency. Secondary adrenal insufficiency is more common than the primary form. It has an estimated prevalence of 150-280 affected per million/year and affects women more frequently than men.

The most common cause of tertiary adrenal insufficiency is the long-term administration of exogenous glucocorticoids, which leads to prolonged suppression of the hypothalamic secretion of the hormone that releases corticotropin.

The main clinical symptoms of adrenal insufficiency include weakness, fatigue, anorexia, nausea and vomiting, abdominal pain, weight loss, postural hypotension and salt craving, while clinical features of a life-threatening adrenal crisis include vomiting, abdominal pain, myalgia, aching joints, severe hypotension and hypovolemic shock.

Before 1949, adrenal insufficiency was a fatal disease. The current availability of cortisone therapy has changed its prognosis. Today, particularly for paediatric patients, extremely effective synthetic products – hydrocortisone and fludrocortisone – are available for the treatment of glucocorticoid and mineralcorticoid deficiencies respectively.

Despite these great therapeutic breakthroughs, the diagnosis and treatment of patients with the disease remains challenging. In particular, fever ($\geq 38^{\circ}\text{C}$), infections, stress, trauma and other acute conditions can drastically alter the metabolic status.

In the event of the onset of “adrenal crisis”, patient management becomes more complex and the prognosis can be poor if the patient is not treated promptly and adequately with increased doses of hydrocortisone, intravenous infusion of physiological saline and glucose.

Main causes of adrenal insufficiency

Tables 1-3 report the main causes of primary, secondary and tertiary adrenal insufficiency (11).

Table 1. Causes of primary adrenal insufficiency

Diseases	Clinical manifestations in addition to adrenal insufficiency
Non genetic disorders	
Autoimmune adrenalitis	
Isolated	Chronic mucocutaneous candidiasis, hypoparathyroidism and other autoimmune diseases (type 1); Thyroid autoimmune disease, type 1 diabetes and other autoimmune diseases (type2); Other autoimmune diseases excluding thyroid disease and type 1 diabetes (type 4)
Syndromic: APS type 1(APECED), type 2 and type 4	
Infectious adrenalitis	
Tuberculous adrenalitis	Symptoms and signs of underlying disease
AIDS	
Fungal adrenalitis	
Infiltrative forms	
Tumours	Symptoms and signs of underlying disease
Non tumors (amyloidosis, haemochromatosis)	
Bilateral adrenal haemorrhage	Meningococcal sepsis, perinatal events
Iatrogenic forms	
Bilateral adrenalectomy	Unresolved Cushing's syndrome, bilateral adrenal masses, pheochromocytoma
Drug-induced adrenal insufficiency (anticoagulants, phenobarbital, ketoconazole, rifampicin, etc.)	None, unless related to drug
Genetic disorders	
Adrenoleukodystrophy	Weakness, spasticity, dementia, blindness, quadriplegia.
Congenital adrenal hyperplasia	
21-hydroxylase deficiency	Hyperandrogenism
11 β -hydroxylase deficiency	Hyperandrogenism, hypertension
Others rarer enzyme deficiencies	
Smith-Lemli-Opitz syndrome	Craniofacial malformations, mental retardation, growth failure, hyponatraemia, hyperkalaemia, cholesterol deficiency
Adrenal hypoplasia congenital	
X-linked	Hypogonadotropic hypogonadism in boys
Xp21 contiguous gene syndrome	Duchenne muscular dystrophy, glycerol kinase deficiency, psychomotor retardation
SF-1-linked	XY sex reversal
Familial glucocorticoid deficiency or corticotropin insensitivity syndromes (type 1, type 2 and variants)	Hyperpigmentation, characteristic facial features, lethargy, muscle, weakness and altered growth rate
Triple A syndrome (Allgrove's syndrome)	Achalasia, alacrima, deafness, mental retardation, hyperkeratosis
Other rare syndromes (IMAGe syndrome, Kearns-Sayre syndrome, Wolman's disease, Sitosterolaemia)	Symptoms and signs of underlying disease

Table 2. Causes of secondary adrenal insufficiency

Diseases	Clinical manifestations in addition to adrenal insufficiency
Space-occupying lesions or trauma	
Pituitary tumors (adenomas, cysts, craniopharyngiomas, ependymomas, meningiomas, rarely carcinomas) or trauma (pituitary stalk lesions)	Panhypopituitarism
Pituitary surgery or irradiation for pituitary tumors, tumors outside the HPA axis or leukemia	Panhypopituitarism
Infections or infiltrative processes	Panhypopituitarism
Pituitary apoplexy	Abrupt onset of severe headache, visual disturbance, nausea, vomiting; panhypopituitarism
Genetic disorders	
Transcription factors involved in pituitary development disorders	Panhypopituitarism and other symptoms related to the specific transcription factor involved
Congenital pro-opiomelanocortin deficiency	Early-onset severe obesity, hyperphagia, red hair
Prader Willi Syndrome	Hypotonia, obesity, mental retardation, hypogonadism

Table 3. Causes of tertiary adrenal insufficiency

Diseases	Clinical manifestations in addition to adrenal insufficiency
Space-occupying lesions or trauma	
Hypothalamic tumours (craniopharyngiomas)	Panhypopituitarism
Hypothalamic surgery or irradiation for CNS or nasopharyngeal tumours	Panhypopituitarism
Infections or infiltrative processes	Panhypopituitarism
Trauma, injury	Panhypopituitarism
Drug-induced adrenal insufficiency	
Glucocorticoid therapy (systemic or topical) or endogenous glucocorticoid hypersecretion (Cushing's syndrome)	Symptoms and signs of underlying disease
Mifepristone	If excessive it can cause severe glucocorticoid deficiency
Antipsychotics(chlorpromazine), antidepressants (imipramine)	None, unless related to drug

Operational guidance

a. Procedures for delivery of therapies

The treatment is provided following the issue of a therapeutic plan by an accredited regional reference centre. As a result of the emergency for the COVID-19 pandemic, AIFA (Agenzia Italiana del Farmaco, the Italian Medicines Agency) has extended the validity to 90 days after expiry of web-based or paper-based treatment plans (PT), signed by specialist doctors (including rare diseases) in March and April (12).

b. Procedures for carrying out clinical tests

Normal practices in use should be adopted. Blood chemistry tests can be performed in any certified analytic laboratory on the advice of the attending physician or of the centre identified by the regions.

c. Management of therapy during infection

In the event of suspected or established COVID-19, patients on replacement therapy must continue the treatment. The dosage of hydrocortisone, which usually varies from 10 to 15 mg/m² of body surface area divided into 2-3 administrations / day depending on the type of adrenal insufficiency, has to be modified as indicated in all cases of acute intercurrent illness, doubling or tripling the dosage, depending on the presence of fever, its duration and on the patient's clinical condition (11-15). If oral intake is impossible (e.g., vomiting), hydrocortisone should be administered parenterally (IM or IV). However, constant contact with the reference center is recommended to allow an assessment by the specialist to prevent adrenal crisis. If a crisis still occurs, urgent hospitalization is necessary. Parents of children with adrenal insufficiency are recommended, as generally suggested by specialists, to make sure they have sufficient quantities of medications, including the intramuscular liquid formulation (IM), even if at present there are no reports of hydrocortisone and fludrocortisone deficiencies.

d. Assessment of infectious risk

Currently there is no epidemiological evidence that both paediatric and adult patients with adrenal insufficiency are at an increased risk of contracting SARS-CoV-2 infection or that the clinical course of the illness would be different from that of the general population. However, patients with adrenal insufficiency must be considered at/as "high risk" in the context of the prevention and management of SARS-CoV-2 infection. They may not be more susceptible to contracting infections than other people, but are at greater risk because febrile episodes can trigger an adrenal crisis, if not properly treated.

e. Preventive measures

Physicians must inform patients with adrenal insufficiency and their parents / guardians, in the case of minors, of the importance of following social distancing measures and of using PPE in case of community life, in order to prevent infection.

The doctor of the territory (Paediatrician / General Practitioner) and the Local Health Service of competence (as currently indicated by the Competent Authority) must be informed if one or more of the

symptoms suggestive of COVID-19 (fever $\geq 38^{\circ}\text{C}$, dry cough, dyspnoea) develop in a patient with adrenal insufficiency. In addition to the symptoms, the particular basic pathology and the followed therapy must be reported. Furthermore, the patient must contact his / her own physician to obtain advice on how to adapt the replacement therapy to the current acute condition unless he/she has already been instructed to increase the dosage independently.

In order to contain the spread of the epidemic during the COVID-19 emergency, routine follow-up visits were deferred in most Italian Paediatric Endocrinology Centres, while retaining the capacity to provide all urgent consultations and the guidance of the treatment in patients with symptoms suggestive of COVID-19.

f. Support of telemedicine systems

The health emergency caused by the COVID-19 pandemic may disrupt routine reference specialist checks. This can occur if the patient is in quarantine or in isolation, or when the rules on social distancing make it impossible for him/her to go to the health facilities for the specialist checks in accordance with the time considered appropriate by the treating specialists.

In some territories, alternative measures have already been activated (e.g. telephone or video calls) to compensate for the reduced capacity of the health system to provide outpatient visits, in order to ensure continuity of care for all paediatric patients with chronic and or rare endocrine diseases.

In some of these clinical events, it is possible to use modern Telemedicine systems to deliver remote medical control services to the patient's home. This can be done by means of video calls, together with the exchange of data and information in digital format that are sent to the doctor directly from the patient or caregiver using software and / or digital medical devices. These technological opportunities can make a significant contribution in the management of continuity of the health care in emergency, if used appropriately within a well-organized care and assistance process in the area. However, the Telemedicine system does not equal Telemedicine technologies, as digital technological innovations alone do not constitute the solution to care problems, being only a means to generate new care strategies and new care services. These methodological aspects are underlined in order to offer healthcare professionals, who deal with patients with adrenal insufficiency, the appropriate scientifically valid points of reference for using these opportunities for the provision of specialist services in an appropriate and safe way.

The definition of the necessary elements for telemedicine services during the COVID-19 emergency is beyond the scope of this work, but more details can be found in the Rapporto ISS COVID-19 n. 12/2020 about interim indications for telemedicine care services during the COVID-19 health emergency (16). Some researchers from COVID-19 Rare Diseases Group also contributed to its implementation.

References

1. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, Tong S. Epidemiology of COVID-19 among children in China. *Pediatrics*. 2020 Mar 16. pii: e20200702. doi: 10.1542/peds.2020-0702.
2. Castagnoli R, Votto M, Licari A, Brambilla I, Bruno R, Perlini S, Rovida F, Baldanti F, Marseglia GL. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review. *JAMA Pediatr*. 2020 Apr 22. doi: 10.1001/jamapediatrics.2020.1467.
3. CDC. Coronavirus disease 2019 in children — United States, February 12–April 2, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:422–426. DOI: <http://dx.doi.org/10.15585/mmwr.mm6914e>.
4. Tagarro A, Epalza C, Santos M, Sanz-Santaefemia FJ, Otheo E, Moraleda C, Calvo C. Screening and Severity of Coronavirus Disease 2019 (COVID-19) in Children in Madrid, Spain. *JAMA Pediatr*. 2020 Apr 8. doi: 10.1001/jamapediatrics.2020.1346.
5. SARS-CoV-2 Surveillance Group. *Characteristics of SARS-CoV-2 patients dying in Italy. Report based on available data on May 7th, 2020*. Roma: Istituto Superiore di Sanità; 2020. https://www.epicentro.iss.it/en/coronavirus/bollettino/Report-COVID-2019_7_may_2020.pdf.
6. Parri N, Lenge M, Buonsenso D; Coronavirus Infection in Pediatric Emergency Departments (CONFIDENCE) Research Group. Children with Covid-19 in pediatric emergency departments in Italy. *N Engl J Med*. 2020 May 1. doi: 10.1056/NEJMc2007617.
7. Zeng H, Xu C, Fan J, Tang Y, Deng Q, Zhang W, Long X. Antibodies in infants born to mothers with COVID-19 pneumonia. *JAMA*. 2020 Mar 26. doi: 10.1001/jama.2020.4861.
8. Stewart MP, Biller MK, Marelli C, Gunnarsson C, Ryan MP, Johannson G. Exploring inpatient hospitalizations and morbidity in patients with adrenal insufficiency. *J Clin Endocrinol Metab* 2016, 101(12), 4843-4850.
9. Bancos I, Hazeldine J, Chortis V, Hampson P, Taylor AE, Lord JM, Arlt W. Primary adrenal insufficiency is associated with impaired natural killer cell function: a potential link to increased mortality. *Eur J Endocrinol* 2017;176(4):471.
10. Tresoldi A, Sumilo D, Perrins M, Toulis KA, Prete A, Narendra Reddy N, Wass JA, Arlt W, Nirantharakumar K. Increased infection risk in Addison's disease and congenital adrenal hyperplasia: a primary care database cohort study. *J Clin Endocrinol Metab* 2020;105(2).
11. Charmandari E, Nicolaidis NC, Chrousos GP. Adrenal insufficiency. *Lancet* 2014 Jun 21;383(9935):2152-67. doi: 10.1016/S0140-6736(13)61684-0.
12. AIFA. *Misure transitorie relative alla proroga dei piani terapeutici AIFA in tema di contenimento e gestione dell'emergenza epidemiologica da Covid-19*. Roma: AIFA; 2020. <https://www.aifa.gov.it/-misure-transitorie-relative-alla-proroga-dei-piani-terapeutici-aifa-in-tema-di-contenimento-e-gestione-dell-emergenza-epidemiologica-da-COVID-19>
13. Kaiser UB, Mirmira RG, Stewart PM. Our response to COVID-19 as endocrinologists and diabetologists. *J Clin Endocrinol Metab*, May 2020, 105(5):1–3. doi:10.1210/clinem/dgaa148.
14. European Society for Paediatric Endocrinology. *COVID-19 information for children and adolescents living with endocrine conditions, including type 1 diabetes mellitus*. Bristol: ESPE; 2020. <https://www.eurospe.org/news/item/14064/COVID-19-information-for-children-and-adolescents-living-with-endocrine-conditions-including-type-1-diabetes-mellitus>
15. European Society of Endocrinology. *COVID-19 and endocrine disease: Clinical information and comment from ESE*. Bristol: ESE; 2020. <https://www.es-e-hormones.org/about-us/our-communities/clinicians/COVID-19-and-endocrine-disease-clinical-information-and-comment-from-ese/>

16. Gabbrielli F, Bertinato L, De Filippis G, Bonomini M, Cipolla M. Indicazioni ad interim per servizi assistenziali di telemedicina durante l'emergenza sanitaria COVID-19. Versione del 13 aprile 2020. Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 12/2020).

Useful links

Società Italiana di Endocrinologia e Diabetologia Pediatrica, SIEDP www.siedp.it

Società Europea di Endocrinologia Pediatrica, ESPE <http://www.eurospe.org>

Società Europea di Endocrinologia ESE www.espe-hormones.org/

Società Italiana di Pediatria, SIP www.sip.it/

WHO COVID-19 www.who.int/emergencies/diseases/novel-coronavirus-2019

SARS-Co V-2 Infection in children: www.nejm.org/doi/full/10.1056/NEJMc2005073

ENDOERN: <https://endo-ern.eu>

Rapporti ISS COVID-19 in Italian

Available from <https://www.iss.it/rapporti-covid-19>

1. Gruppo di lavoro ISS Prevenzione e controllo delle Infezioni.
Indicazioni ad interim per l'effettuazione dell'isolamento e della assistenza sanitaria domiciliare nell'attuale contesto COVID-19. Versione del 7 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020 (Rapporto ISS COVID-19, n. 1/2020)
2. Gruppo di lavoro ISS Prevenzione e controllo delle Infezioni.
Indicazioni ad interim per un utilizzo razionale delle protezioni per infezione da SARS-CoV-2 nelle attività sanitarie e sociosanitarie (assistenza a soggetti affetti da COVID-19) nell'attuale scenario emergenziale SARS-CoV-2. Versione del 28 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020 (Rapporto ISS COVID-19, n. 2 Rev./2020)
3. Gruppo di lavoro ISS Ambiente e Gestione dei Rifiuti.
Indicazioni ad interim per la gestione dei rifiuti urbani in relazione alla trasmissione dell'infezione da virus SARS-CoV-2. Versione del 31 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020 (Rapporto ISS COVID-19, n. 3 Rev./2020)
4. Gruppo di lavoro ISS Prevenzione e controllo delle Infezioni.
Indicazioni ad interim per la prevenzione e il controllo dell'infezione da SARS-CoV-2 in strutture residenziali sociosanitarie. Versione del 17 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020 (Rapporto ISS COVID-19, n. 4/2020 Rev.)
5. Gruppo di lavoro ISS Ambiente e Qualità dell'aria indoor.
Indicazioni ad interim per la prevenzione e gestione degli ambienti indoor in relazione alla trasmissione dell'infezione da virus SARS-CoV-2. Versione del 21 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 5/2020 Rev.)
6. Gruppo di lavoro ISS Cause di morte COVID-19.
Procedura per l'esecuzione di riscontri diagnostici in pazienti deceduti con infezione da SARS-CoV-2. Versione del 23 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 6/2020).
7. Gruppo di lavoro ISS Biocidi COVID-19 e Gruppo di lavoro ISS Ambiente e Rifiuti COVID-19.
Raccomandazioni per la disinfezione di ambienti esterni e superfici stradali per la prevenzione della trasmissione dell'infezione da SARS-CoV-2. Versione del 29 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 7/2020).
8. Osservatorio Nazionale Autismo ISS.
Indicazioni ad interim per un appropriato sostegno delle persone nello spettro autistico nell'attuale scenario emergenziale SARS-CoV-2. Versione del 30 marzo 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 8/2020).
9. Gruppo di Lavoro ISS Ambiente – Rifiuti COVID-19.
Indicazioni ad interim sulla gestione dei fanghi di depurazione per la prevenzione della diffusione del virus SARS-CoV-2. Versione del 3 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 9/2020).
10. Gruppo di Lavoro ISS Ambiente-Rifiuti COVID-19.
Indicazioni ad interim su acqua e servizi igienici in relazione alla diffusione del virus SARS-CoV-2 Versione del 7 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 10/2020).

11. Gruppo di Lavoro ISS Diagnostica e sorveglianza microbiologica COVID-19: aspetti di analisi molecolare e sierologica
Raccomandazioni per il corretto prelievo, conservazione e analisi sul tampone oro/nasofaringeo per la diagnosi di COVID-19. Versione del 7 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 11/2020).
12. Gabbrielli F, Bertinato L, De Filippis G, Bonomini M, Cipolla M.
Indicazioni ad interim per servizi assistenziali di telemedicina durante l'emergenza sanitaria COVID-19. Versione del 13 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 12/2020).
13. Gruppo di lavoro ISS Ricerca traslazionale COVID-19.
Raccomandazioni per raccolta, trasporto e conservazione di campioni biologici COVID-19. Versione del 15 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 13/2020).
14. Gruppo di lavoro ISS Malattie Rare COVID-19.
Indicazioni ad interim per un appropriato sostegno delle persone con enzimopenia G6PD (favismo) nell'attuale scenario emergenziale SARS-CoV-2. Versione del 14 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 14/2020).
15. Gruppo di lavoro ISS Farmaci COVID-19.
Indicazioni relative ai rischi di acquisto online di farmaci per la prevenzione e terapia dell'infezione COVID-19 e alla diffusione sui social network di informazioni false sulle terapie. Versione del 16 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 15/2020).
16. Gruppo di lavoro ISS Sanità Pubblica Veterinaria e Sicurezza Alimentare COVID-19.
Animali da compagnia e SARS-CoV-2: cosa occorre sapere, come occorre comportarsi. Versione del 19 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 16/2020).
17. Gruppo di lavoro ISS Sanità Pubblica Veterinaria e Sicurezza Alimentare COVID-19.
Indicazioni ad interim sull'igiene degli alimenti durante l'epidemia da virus SARS-CoV-2. Versione del 19 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 17/2020).
18. Gruppo di lavoro ISS Ricerca traslazionale COVID-19.
Raccomandazioni per la raccolta e analisi dei dati disaggregati per sesso relativi a incidenza, manifestazioni, risposta alle terapie e outcome dei pazienti COVID-19. Versione del 26 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 18/2020).
19. Gruppo di lavoro ISS Biocidi COVID-19.
Raccomandazioni ad interim sui disinfettanti nell'attuale emergenza COVID-19: presidi medico-chirurgici e biocidi. Versione del 25 aprile 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 19/2020).
20. Gruppo di Lavoro ISS Prevenzione e Controllo delle Infezioni.
Indicazioni per la sanificazione degli ambienti interni per prevenire la trasmissione di SARS-COV 2. Versione dell'8 maggio 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 20/2020).
21. Ricci ML, Rota MC, Scaturro M, Veschetti E, Lucentini L, Bonadonna L, La Mura S.
Guida per la prevenzione della contaminazione da Legionella negli impianti idrici di strutture turistico recettive e altri edifici ad uso civile e industriale, non utilizzati durante la pandemia COVID-19. Versione del 3 maggio 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 21/2020).

22. Gruppo di lavoro ISS Salute mentale ed emergenza COVID-19
Indicazioni ad interim per la gestione dello stress lavoro-correlato negli operatori sanitari e socio-sanitari durante lo scenario emergenziale SARS-COV-2. Versione del 7 maggio.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 22/2020)

23. Gruppo di lavoro ISS Salute mentale ed emergenza COVID-19
Indicazioni di un programma di intervento dei Dipartimenti di Salute Mentale per la gestione dell'impatto dell'epidemia COVID-19 sulla salute mentale. Versione del 6 maggio 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 23/2020).

24. Gruppo di lavoro ISS Malattie Rare COVID-19.
Indicazioni ad interim per una appropriata gestione dell'ipossurrenalismo in età pediatrica nell'attuale scenario emergenziale da infezione da SARS-CoV-2. Versione del 10 maggio 2020.
Roma: Istituto Superiore di Sanità; 2020. (Rapporto ISS COVID-19, n. 24/2020)