Articles

Long-term survival for lymphoid neoplasms and national health expenditure (EUROCARE-6): a retrospective, population-based study



Milena Sant, Claudia Vener, Roberto Lillini, Silvia Rossi, Simone Bonfarnuzzo, Rafael Marcos-Gragera, Marc Maynadié, Kaire Innos, Keiu Paapsi, Otto Visser, Alice Bernasconi, Elena Demuru, Corrado Di Benedetto, Seyed Mohsen Mousavi, Marcel Blum, Philip Went, Diego Serraino, Damien Bennett, Maria-Jose Sánchez, Roberta De Angelis, and the EUROCARE-6 Working Group*

Summary

Background Management of lymphoid malignancies requires substantial health system resources. Total national health expenditure might influence population-based lymphoid malignancy survival. We studied the long-term survival of patients with 12 lymphoid malignancy types and examined whether different levels of national health expenditure might explain differences in lymphoid malignancy prognosis between European countries and regions.

Methods For this observational, retrospective, population-based study, we analysed the EUROCARE-6 dataset of patients aged 15 or older diagnosed between 2001 and 2013 with one of 12 lymphoid malignancies defined according to International Classification of Disease for Oncology (third edition) and WHO classification, and followed up to 2014 (Jan 1, 2001–Dec 31, 2014). Countries were classified according to their mean total national health expenditure quartile in 2001–13. For each lymphoid malignancy, 5-year and 10-year age-standardised relative survival (ASRS) was calculated using the period approach. Generalised linear models indicated the effects of age at diagnosis, gender, and total national health expenditure on the relative excess risk of death (RER).

Findings 82 cancer registries (61 regional and 21 national) from 27 European countries provided data eligible for 10-year survival estimates comprising 890730 lymphoid malignancy cases diagnosed in 2001–13. Median follow-up time was 13 years (IQR 13–14). Of the 12 lymphoid malignancies, the 10-year ASRS in Europe was highest for hairy cell leukaemia (82.6% [95% CI 78.9–86.5) and Hodgkin lymphoma (79.3% [78.6–79.9]) and lowest for plasma cell neoplasms (29.5% [28.9–30.0]). RER increased with age at diagnosis, particularly from 55–64 years to 75 years or older, for all lymphoid malignancies. Women had higher ASRS than men for all lymphoid malignancies, except for precursor B, T, or natural killer cell, or not-otherwise specified lymphoblastic lymphoma or leukaemia. 10-year ASRS for each lymphoid malignancy was higher (and the RER lower) in countries in the highest national health expenditure quartile than in countries in the lowest quartile, with a decreasing pattern through quartiles for many lymphoid malignancies. 10-year ASRS for non-Hodgkin lymphoma, the most representative class for lymphoid malignancies based on the number of incident cases, was 59.3% (95% CI 58.7-60.0) in the first quartile, 57.6% (55.2-58.7) in the second quartile, 55.4% (54.3-56.5) in the third quartile, and 44.7% (43.6-45.8) in the fourth quartile; with reference to the European mean, the RER was 0.80 (95% CI 0.79-0.82) in the first, 0.91 (0.90-0.93) in the second, 0.94 (0.92-0.96) in the third, and 1.45 (1.42-1.48) in the fourth quartiles.

Interpretation Total national health expenditure is associated with geographical inequalities in lymphoid malignancy prognosis. Policy decisions on allocating economic resources and implementing evidence-based models of care are needed to reduce these differences.

Funding Italian Ministry of Health, European Commission, Estonian Research Council.

Copyright © 2024 Elsevier Ltd. All rights reserved, including those for text and data mining, AI training, and similar technologies.

Introduction

Haematological malignancies account for approximately 7% of all cancers worldwide¹ and more than half are lymphoid malignancies. In 2020, the agestandardised incidence rates (per 100 000) in Europe for both genders were estimated as 16.4 for non-Hodgkin lymphoma, 6.8 for multiple myeloma, and 2.7 for Hodgkin lymphoma.² The incidence of lymphoid malignancies is higher in men than in women and increases with age,² typically affecting older people.³

The improvements in survival reported by the EUROCARE-5 study for many lymphoid malignancies coincided with the introduction of new treatments, such as anti-CD20 monoclonal antibodies for treating B-cell lymphomas and acute lymphoblastic leukaemia or proteasome inhibitors for multiple myeloma.⁴ The development of chimeric antigen receptor redirected

Lancet Oncol 2024; 25: 731–43

Published Online May 1, 2024 https://doi.org/10.1016/ S1470-2045(24)00141-4

See Comment page 684

*For the members of the EUROCARE-6 Working Group see the appendix (pp 26–27) Analytical Epidemiology and

Health Impact Unit (M Sant MD. C Vener MD, R Lillini PhD, S Bonfarnuzzo), Evaluative **Epidemiology Unit** (A Bernasconi MSc). and **Epidemiology and Prevention** Unit (C Vener), Department of Epidemiology and Data Science, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy: Department of **Oncology and Molecular** Medicine, Istituto Superiore di Sanità, Rome, Italy (S Rossi MSc, E Demuru PhD, C Di Benedetto MSc. R De Angelis MSc); Epidemiology Unit and Girona Cancer Registry, Oncology Coordination Plan, Department of Health Government of Catalonia. Catalan Institute of Oncology, Girona, Spain (R Marcos-Gragera MD); **Registre des Hémopathies** Malignes de Côte d'Or. University of Burgundy and Dijon University Hospital, Diion, France (Prof M Maynadié MD); Department of Epidemiology and Biostatistics, National Institute for Health Development, Tallinn, Estonia (K Innos PhD, K Paapsi MSc): Department of Registration, **Netherlands Comprehensive** Cancer Organization, Utrecht, Netherlands (O Visser MD); Cancer Registry Graubünden-Glarus, Chur, Switzerland (S M Mousavi MD); East Switzerland Cancer Registry and Cancer Registry Liechtenstein, St Gallen, Switzerland (M Blum MA);

Institut für Pathologie, Kantonsspital Graubünden, Chur, Switzerland (P Went MD); Cancer Registry of Friuli-Venezia Giulia, Aviano, Italy (D Serraino MD); Northern Ireland Cancer Registry (NICR), Belfast, UK (D Bennett PhD); **Oueens University Belfast** (QUB), Centre for Public Health, Institute of Clinical Sciences Royal Victoria Hospital, Belfast, UK (D Bennett); Escuela Andaluza de Salud Pública (EASP), Granada, Spain (M-J Sánchez MD); Instituto de Investigación Biosanitaria ibs Granada, Granada, Spain (M-J Sánchez); Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain (M-J Sánchez); Department of Preventive Medicine and Public Health, University of Granada, Granada, Spain (M-J Sánchez)

Correspondence to: Dr Claudia Vener, Epidemiology and Prevention Unit, Department of Epidemiology and Data Science, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, 20133 Milan, Italy

claudia.vener@istitutotumori. mi.it

See Online for appendix

Research in context

Evidence before this study

Cancer registry-based survival studies on lymphoid malignancies usually provide 5-year survival estimates, whereas long-term survival is available for specific malignancy types or on a national basis only. The improvements in 5-year survival reported by previous EUROCARE studies for many lymphoid malignancies coincided with the introduction of new treatments but persisting survival differences across Europe suggest inequalities in quality of care and access to treatments exist. Total national expenditure on health—an indicator of a country's overall investment in its health system—was associated with 5-year all-cancer survival, but its influence on long-term survival for specific lymphoid malignancies has not been investigated. No formal literature review was done before the start of the study.

Added value of this study

To our knowledge, this is the first study to provide 10-year survival estimates for 12 specific lymphoid malignancies in 27 European countries, using data from 82 population-based cancer registries. It is also the first study to systematically analyse the relationship between survival for each lymphoid malignancy and total national health expenditure. For all the lymphoid malignancies, 5-year and 10-year age-standardised relative survival (ASRS) in countries in the highest total national health expenditure quartile was higher than in countries in the lowest quartile, with a gradual decrease in survival through decreasing expenditure quartiles, although this pattern differed by lymphoid malignancy subgroup. For common lymphoid malignancies, such as Hodgkin lymphoma and non-Hodgkin lymphoma, survival was similar across the countries in the high and medium-high quartiles. 10-year ASRS for Hodgkin lymphoma decreased from the highest to lowest quartile. The decrease through quartiles was less uniform for lymphoid malignancies with poor prognosis, such as mantle cell lymphoma. In most countries in the first or second total national health expenditure quartiles, 10-year ASRS was in line with or above the European mean for common lymphoid malignancies. Exceptions were England, Scotland, Northern Ireland, Finland, and Austria, where survival for some lymphoid malignancies was lower than the European mean. Multivariable survival models (adjusted by age, gender, and total national health expenditure), applied for each lymphoid malignancy, confirmed that 10-year survival was better in countries with higher healthcare expenditure than in those with lower expenditures.

Implications of all the available evidence

In countries with low total national health expenditure, policy decisions are needed to increase health-care resources and identify strategies to sustain interventions. Differences in lymphoid malignancy survival among countries with similar total national health expenditure indicate the need to investigate national health-care policies and health resource utilisation. All these countries need to implement efficient, evidence-based care and long-term surveillance to reduce inequalities. Given the high cost of drugs for haematological malignancies, political pressure and negotiations with drug manufacturers to reduce the prices of the most expensive drugs should be pursued.

T-cell (CAR-T) therapy⁵ and other therapeutic targets (eg, BTK inhibitors) is expected to improve survival for patients with various B-cell neoplasms.⁶ However, persistent survival inequalities across Europe suggest differences in quality of care and access to treatments exist.⁷

The high and increasing costs of lymphoid malignancy management—due to expensive drugs, complex regimens, and the need for extensive support services and after-care surveillance—require substantial health system resources.⁸ Total national expenditure on health was associated with 5-year all-cancer survival in the EUROCARE-5 study,⁹ but its association with long-term survival for specific lymphoid malignancies has not been investigated.

As increasing numbers of patients with haematological malignancies in Europe survive for more than 5 years after their initial diagnosis,^{10,11} survival needs to be analysed beyond the conventional 5-year timeframe usually reported. Studies investigating long-term survival indicated that follicular lymphoma^{12,13} survival continues to decrease over the follow-up years and mortality remained higher than in the general population even

10 years after diagnosis.¹⁰ Studies in children and adolescents reported that the persisting excess risk of death¹⁴ might be attributable to secondary cancers, late relapses of non-Hodgkin lymphoma, or cardiovascular disease.¹⁵

Apart from late relapses, the excess mortality of patients with lymphoid malignancies, persisting for more than 5 years after diagnosis, has also been attributed to the long-term adverse effects of cytotoxic therapy,¹⁶ multimorbidity,¹⁷ and social deprivation.¹⁸

Total national health expenditure represents the country's general resources for health, allowing us to evaluate how the economic availabilities can contribute to differences in health outcomes and permitting reliable results comparison among countries. It is a sufficiently standardised macroeconomic indicator collected by national and international statistics offices. Total national health expenditure has already been considered for evaluating differences in non-Hodgkin lymphoma survival across countries.¹⁹ It is a general descriptor that is useful to uncover potential issues to be better explored by future studies, outside the EUROCARE-6 framework.

We studied the long-term survival of patients with 12 specific lymphoid malignancy types and examined whether different levels of total national health expenditure might explain differences in lymphoid malignancy prognosis between European countries and regions.

Methods

Study design and data collection

For this observational, retrospective, population-based study, we selected 10-year survival data of adults (aged \geq 15 years) diagnosed with lymphoid malignancies from the dataset of the EUROCARE-6 study, which relies on standardised information provided by population-based cancer registries in Europe. The quality and completeness of cancer registry data were assessed with standardised EUROCARE and European Network of Cancer Registries and Joint Research Centre checking procedures.²⁰ The study protocol is available online.

Morphology was coded according to the International Classification of Disease for Oncology, third edition (ICDO-3),²¹ and 2017 WHO classification (WHO-4),²² with some updates introduced in the 2023 revision. We collected data on date of birth, diagnosis, and death, vital status, and gender through patients' medical records held at cancer registries.

The lymphoid malignancy types analysed were Hodgkin lymphoma, the group of non-Hodgkin lymphoma, and ten specific types: chronic lymphocytic leukaemia/small lymphocytic lymphoma; lymphoplasmacytic lymphoma/ Waldenström macroglobulinaemia; diffuse large B-cell lymphoma; mantle cell lymphoma; follicular lymphoma; marginal zone lymphoma; mature T and natural killer cell lymphoma; precursor B-cell, T-cell, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia; plasma cell neoplasms; and hairy cell leukaemia. Detailed ICDO morphology codes of the lymphoid malignancy forms analysed with their acronyms are provided in the appendix (p 13).

Statistical analysis

We computed relative survival (the cancer-specific survival probability) with the Ederer II method, using SEER*Stat software and registry-specific lifetables stratified by age, gender, and year. 5-year and 10-year relative survival in the follow-up period (2010-14; selecting patients diagnosed in 2001-13), with 95% CIs, were calculated by country, gender, and age at diagnosis using the period approach, which provides the most upto-date estimates of the survival of patients diagnosed in 2010-14.23 In our analyses, the Ederer II estimator was less prone than the net survival estimator to estimates' instabilities or increases for long follow-up durations that typically occur with scarce number of cases and in older age groups.²⁴ Relative survival SEs were calculated with Greenwood's formula assuming fixed expected survival. Two-sided 95% CIs were computed using logarithmic transformation.

Age at diagnosis was divided into five groups (15–44, 45–54, 55–64, 65–74, and ≥75 years) and reduced to four for Hodgkin lymphoma and precursor B-cell, T-cell, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia (15–44, 45–54, 55–64, and ≥65 years) and for mantle cell lymphoma and marginal zone lymphoma (15–54, 55–64, 65–74, and ≥75 years).

To account for different age structures of European populations, country-specific relative survival estimates were standardised by age with international cancer survival standards.²⁵ European mean survival was computed by weighting area-specific survival by the population of each area (northern, central, southern, and eastern Europe; UK; and Ireland).

The total national health expenditure—in US\$ purchasing power parity per capita—by country was determined annually for 2001–13 from key international databases (eg, Health for All Europe, Organisation for Economic Co-operation and Development, World Bank Database, and local national statistics offices).

Countries were scored according to their mean total national health expenditure in 2001–13 and divided into quartiles of expenditure (low, middle-low, middle-high, and high). We adopted total national health expenditure quartiles for easier interpretation of results, while accepting the loss of some precision and leaving for future analyses the application of more structured methods (ie, fractional polynomial or regression spline scales). Reliability and standardisation of total national health expenditure data made this choice reasonable.

The *I*² statistic was calculated to establish the percentage of total between-country survival differences due to heterogeneity rather than chance.²⁶ The higher the *I*² statistic, the more the differences are due to true heterogeneity between the populations.

Generalised linear models were used in multivariable regressions to assess the effects of age at diagnosis, gender, and total national health expenditure on the relative excess risks of death (RERs), with 95% CIs, for each lymphoid malignancy group in the 10 years after diagnosis. The selection of the variables was determined by their availability in cancer registry databases. We used a Poisson regression model, in which the response variable was the difference between observed and estimated deaths and the predictors were age at diagnosis (grouped as described earlier), gender, and total national health expenditure in quartiles (grouped as above; appendix p 26).²⁷

The possible combined effects of age, gender, and total national health expenditure on RERs were investigated. After considering only the main effects, generalised linear models with both two-way and three-way interaction effects along with the main effects were computed.²⁸ Given the high number of cases, Hodgkin lymphoma and non-Hodgkin lymphoma groups were considered for examining possible association between total national health expenditure and RERs in subgroups

For more on the **EUROCARE-6 data** see https://encr.eu/sites/ default/files/Data_call/2015_ ENCR_JRC_Call_for_Data_ Version_1_1.pdf

For the **study protocol** see https://www.iss.it/en/eurocare-6

For more on the **morphology coding** see https://seer.cancer. gov/tools/heme/

For the health data from the Organisation for Economic Co-operation and Development see https://www. oecd.org/els/health-systems/ health-data.htm

For the **SEER*Stat software** see https://seer.cancer.gov/seerstat/

defined by age (15–54 and 55+ years) and gender.²⁹ The computation of RERs by generalised linear model were repeated for these subgroups.

Data were analysed using SEER*Stat software (version 8.3.9), SAS (version 9.4), and STATA (version 17.0).

Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

Of 913 244 eligible lymphoid malignancy cases diagnosed in 27 European countries between 2001 and 2013 (table 1),

1956 (0.2%) were excluded because of major errors (missing or invalid data and inconsistencies in essential variables to compute survival, such as dates, information on vital status, basis of diagnosis, topography, and morphology). Moreover, 16 921 (1.9%) cases known only from death certificates and 2396 (0.3%) incidentally detected at autopsy were also excluded. The proportions of cases known only from death certificates were relatively high in Germany, Austria, Bulgaria, and Slovakia (table 1).

The number of cases alive with unknown survival time was 1239 (0.1%); the number of cases lost to follow-up (ie, censored alive before the follow-up closing date) 624 (0.5%) at 5 years, with the highest proportions in

	All lymphoid malignancy cases, 2001–13	Invalid cases e	xcluded from sur	vival analysis, n	(%)	Data quality i	ndicator, n (%)
		Major errors	Death certificate only	Incidentally detected at autopsy	Alive cases with unknown survival time	Lost to follow-up*	NOS morphology†
Austria	27330	0	1703 (6.2%)	0	0	0	4746 (18·5%)
Bulgaria	13840	24 (0.2%)	730 (5·3%)	0	0	0	2514 (19·2%)
Croatia	11877	0	0	0	78 (0.7%)	0	3230 (27.4%)
Czechia	34289	20 (0.1%)	281 (0.8%)	1546 (4·5%)	172 (0.5%)	0	4770 (14·8%)
Denmark	24363	36 (0.1%)	0	35 (0.1%)	2 (<0.1%)	15 (0.5%)	1588 (6.5%)
Estonia	4035	0	61 (1.5%)	50 (1·2%)	0	4 (0.9%)	297 (7.6%)
Finland	24465	614 (2.5%)	2 (<0.1%)	13 (0.1%)	0	0	4324 (18·1%)
France (13 cancer registries)	53792	0	0	0	101 (0.2%)	83 (1.0%)	2503 (4.7%)
Germany (5 cancer registries)	76666	41 (0.1%)	6334 (8·3%)	23 (<0.1%)	111 (0.1%)	77 (0.9%)	6181 (8.8%)
Iceland	1128	0	3 (0.3%)	7 (0.6%)	0	0	30 (2.7%)
Ireland	14649	1(<0.1%)	73 (0.5%)	58 (0.4%)	0	0	2210 (15.2%)
Italy (29 cancer registries)	93602	10 (<0.1%)	484 (0·5%)	136 (0.1%)	199 (0·2%)	290 (1·9%)	10583 (11·4%)
Latvia	5153	7 (0.1%)	1(<0.1%)	0	0	0	587 (11·4%)
Lithuania	9295	0	54 (0.6%)	6 (0.1%)	11 (0.1%)	15 (1·4%)	653 (7.1%)
Malta	1341	2 (0.1%)	1(0.1%)	21 (1.6%)	0	0	194 (14·7%)
Netherlands	68 531	514 (0.8%)	0	203 (0.3%)	0	0	1386 (2.0%)
Norway	21934	294 (1·3%)	120 (0·5%)	5 (<0.1%)	0	0	994 (4.6%)
Poland	81286	90 (0·1%)	1457 (1·8%)	48 (0.1%)	475 (0.6%)	0	24340 (30·7%)
Portugal (2 cancer registries)	23245	56 (0.2%)	8 (<0.1%)	0	58 (0.2%)	44 (1·2%)	5342 (23.1%)
Slovakia	11753	4 (<0.1%)	605 (5·1%)	46 (0.4%)	0	0	399 (3.6%)
Slovenia	6556	0	0	83 (1.3%)	1(<0.1%)	0	283 (4.4%)
Spain (8 cancer registries)	29115	116 (0.4%)	244 (0.8%)	54 (0·2%)	5 (<0.1%)	17 (0.3%)	1414 (4.9%)
Switzerland (4 cancer registries)	7196	0	27 (0.4%)	24 (0·3%)	26 (0.4%)	71 (5.8%)	297 (4·2%)
England (UK)	223285	93 (<0.1%)	4339 (1·9%)	0	0	0	26 471 (12.1%)
Northern Ireland (UK)	7028	2 (<0.1%)	46 (0.7%)	8 (0.1%)	0	0	1013 (14.5%)
Scotland (UK)	24250	2 (<0.1%)	62 (0.3%)	32 (0.1%)	0	8 (0.3%)	2042 (8.5%)
Wales (UK)	13240	28 (0.2%)	286 (2.2%)	0	0	0	1461 (11·3%)
European pool (82 cancer registries)	913244	1956 (0.2%)	16921(1.9%)	2396 (0.3%)	1239 (0.1%)	624 (0.5%)	109 852 (12·3%)

Data quality indicators are shown by country, with numbers of cases by specific malignancy. NOS=not otherwise specified. *Proportion of patients diagnosed in 2005–08, censored on Dec 31, 2013, with less than 5 years of follow-up; the proportion is calculated for cases diagnosed in 2005–07 in Croatia and Germany (two of five cancer registries), where follow-up closing date was Dec 31, 2012. †NOS International Classification of Disease for Oncology, third edition, morphology for lymphoid malignancies: 9590, 9591, 9820, 9832.

Table 1: Data checks for adults (aged ≥15 years) diagnosed with lymphoid malignancy in 2001–13 in Europe

	Alllymphoid malignancies	Hodgkin lymphoma	Non- Hodgkin lymphoma	Chronic Iymphocytic leukaemia/ small Iymphocytic Iymphoma	Lymphoplasmacytic lymphoma/ Waldenström macroglobulinaemia	Diffuselarge B-cell lymphoma	Mantle cell lymphoma	Follicular lymphoma	Marginal zone lymphoma	Mature T-cell and natural killer cell lymphoma	Precursor B-cell, T-cell, natural killer cell, or not- otherwise- specified lymphoblastic lymphoma/ leukaemia	Plasma cell neoplasms	Hairy cell leukaemia
Austria	25627	1981	13 987	4478	567	3066	655	1956	926	1312	579	4748	330
Bulgaria	13086	1868	6064	2541	187	1789	159	478	43	432	623	2187	184
Croatia	11797	1293	5502	2221	108	864	72	875	73	137	415	2305	129
Czechia	32 270	3207	16842	6864	366	5326	816	2495	663	1338	661	5099	294
Denmark	24290	1671	12688	5275	207	4640	666	2409	752	951	454	4425	253
Estonia	3924	390	1761	1040	49	751	113	119	94	130	107	739	40
Finland	23 836	1724	13 911	3791	NA	NA	NA	NA	NA	NA	443	4526	210
France (13 cancer registries)	53 691	3946	29193	10870	2966	8925	1392	4840	3205	3041	1039	9663	586
Germany (5 cancer registries)	70157	4905	36125	13756	1731	11697	1953	6479	2897	3530	1346	14212	674
Iceland	1118	108	572	187	46	199	25	123	56	58	29	233	∞
Ireland	14517	1276	7524	2775	273	2163	245	1291	249	591	331	2786	175
Italy (29 cancer registries)	92 773	8260	50 418	13742	2494	15716	1821	6767	4515	4525	2217	20142	1038
Latvia	5145	586	2104	1380	55	504	15	172	477	222	178	878	26
Lithuania	9224	862	3954	2633	108	1486	231	196	326	544	274	1749	72
Malta	1317	149	833	100	16	256	17	111	38	162	49	206	12
Netherlands	67814	5056	37983	11570	2546	15105	2169	6748	3183	3421	1372	13 446	782
Norway	21515	1514	11592	3358	1034	4103	561	2333	892	1070	429	4660	216
Poland	79216	9855	33 615	17 513	540	279	5	3928	1165	1955	2441	16194	587
Portugal (2 cancer registries)	23123	2386	14055	2414	336	3505	359	1808	782	1079	486	4148	176
Slovakia	11098	1230	4642	2878	176	2016	214	662	229	379	355	2348	113
Slovenia	6472	539	3331	1158	128	1373	230	416	444	293	155	1304	58
Spain (8 cancer registries)	28 696	2751	15655	4579	649	5288	660	3304	1612	1676	694	5469	200
Switzerland (4 cancer registries)	7119	654	3901	1295	209	1345	242	677	427	371	128	1277	93
England (UK)	218 853	16411	117 442	36358	5840	39458	4655	20 217	6541	7998	3785	47 174	2078
Northern Ireland (UK)	6972	620	3805	1001	109	1096	83	837	144	278	115	1567	66
Scotland (UK)	24 154	1889	12 938	4022	691	4725	447	2472	925	1122	399	4972	169
Wales (UK)	12 926	854	6836	2532	310	2402	286	1147	401	430	195	2700	119
European pool (82 cancer registries)	890730	75 985	467 273	160331	22 441	138 077	18 091	72 860	31 059	37 045	19299	179 157	8688
NA=not available.													
Table 2: Number of a	dult (aged ≥15 ye	ears) cases of ly	ymphoid mali	gnancy, by speci	fic malignancy, diagnos	ed in 2001–13	in Europe, incl	luded in survi	val analysis				



Figure 1: 10-year age-standardised relative survival for adult patients followed up in 2010–14 in Europe for each lymphoid malignancy

Data are presented as percentages with 95% CIs indicated by error bars.

Switzerland and Italy (table 1). Cases lost to follow-up were not excluded from the analysis; they contributed for the whole period in which they were under observation, up to their censoring date. The percentages of nototherwise-specified morphologies were higher in Poland, Croatia, Portugal, Bulgaria, Austria, and Finland than the European average (table 1).

Following these checks and exclusions. 890730 lymphoid malignancy cases diagnosed in 2001-13 were included in the survival analysis (table 2). Results from the quality checks of cancer registry-specific data are shown in the appendix (pp 14-16). Of the 100 European cancer registries that submitted data, 82 registries (61 regional and 21 national) from 27 European countries provided data eligible for 10-year survival estimates-ie, data on adult cases diagnosed with lymphoid malignancies in 2001-13 and followed up until the end of 2014 (Jan 1, 2001, to Dec 31, 2014). Median follow-up time was 13 years (IQR 13-14). The categorisation of the countries on the basis of their total national health expenditure is shown in the appendix (pp 24-25).

Of the 12 lymphoid malignancies, the 10-year agestandardised relative survival (ASRS) in Europe was highest for hairy cell leukaemia (82.6% [95% CI 78.9-86.5]) and Hodgkin lymphoma (79.3% [78.6-79.9]) and lowest for plasma cell neoplasms (29.5% [28.9-30.0]). 10-year survival was similar for chronic lymphocytic leukaemia/small lymphocytic lymphoma (56.8%[56.1-57.5]) and non-Hodgkin lymphoma (55.3% [54·9–55·8]) with higher survival for the non-Hodgkin lymphoma types: follicular lymphoma (71·0% [69·6–72·4]) and marginal zone lymphoma (70·1% [68·4–71·9%]); survival was lower for diffuse large B-cell lymphoma (49·9% [49·2–50·7]) and mature T and natural killer cell lymphoma (49·9% [48·5–51·3]) and considerably lower for mantle cell lymphoma (37·9% [36·1–39·8]); lymphoplasmacytic lymphoma/Waldenström macroglobulinaemia had an intermediate prognosis (59·4% [57·4–61·4]). Finally, precursor B, T, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia showed a poor prognosis (41·4% [39·9–42·9]; appendix pp 17–20).

Women generally had a better prognosis than men; the largest gender differences (5–6 percentage points) were for the whole non-Hodgkin lymphoma group and for the subtypes of lymphoplasmacytic lymphoma/Waldenström macroglobulinaemia, chronic lymphocytic leukaemia/ small lymphocytic lymphoma, mantle cell lymphoma, and marginal zone lymphoma (figure 1).

For all lymphoid malignancies, relative survival decreased with older age (figure 2; appendix pp 2–7). For Hodgkin lymphoma, 10-year relative survival decreased from $92 \cdot 3\%$ (95% CI $91 \cdot 8 - 92 \cdot 8$) in the youngest age class to $42 \cdot 8\%$ ($40 \cdot 4 - 45 \cdot 4$) in the oldest. 10-year relative survival was lowest for patients with precursor B, T, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia aged 65 years or older ($11 \cdot 8\%$ [95% CI $9 \cdot 7 - 14 \cdot 3$] compared with 56 $\cdot 1\%$ [53 $\cdot 9 - 58 \cdot 4$] for the youngest group). For all lymphoid malignancies, age-specific relative survival was higher for women than men (data not shown).

The I^2 statistic (measuring heterogeneity not due to chance) was lowest for precursor B, T, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia (73.5%) and highest for non-Hodgkin lymphoma (98.2%), chronic lymphocytic leukaemia/ small lymphocytic lymphoma (97.6%), and plasma cell neoplasms (93.8%; figure 3). For all other lymphoid malignancies, the I^2 exceeded 80%.

The largest European mean ASRS decrease from the 5th to the 10th year after diagnosis was around 16–19% for chronic lymphocytic leukaemia/small lymphocytic lymphoma, lymphoplasmacytic lymphoma, Waldenström macroglobulinaemia, mantle cell lymphoma, and plasma cell neoplasms (appendix pp 17–20). There were smaller decreases (8–10%) for diffuse large B-cell lymphoma, follicular lymphoma, and marginal zone lymphoma, and for the whole non-Hodgkin lymphoma group. There were less marked differences (around 5–6%) for Hodgkin lymphoma, mature T and natural killer cell lymphoma, precursor B, T, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia, and hairy cell leukaemia.

For all the lymphoid malignancies, 5-year and 10-year ASRS in countries in the highest total national health expenditure quartile (high expenditure) was higher than



Figure 2: 10-year relative survival for adult patients followed up in 2010–14 in Europe, by age group for each lymphoid malignancy Data are presented as percentages with 95% CIs in parentheses.

in countries in the lowest quartile (low expenditure), with a graduated decrease in survival through the expenditure quartiles, although this pattern varied by lymphoid malignancy subgroup. The extent of these differences was similar for 5-year and 10-year ASRS. Survival was similar and decreased only slightly from the first to the fourth total national health expenditure quartile for Hodgkin lymphoma, showing 10-year relative survival values of 82.7% (95% CI 81.6-83.8) in the first, 80.7% (79.8–81.6) in the second, 76.1%



Figure 3: Across-country variation in 10-year age-standardised relative survival, by each lymphoid malignancy

Outliers fell outside the whiskers' length, defined as 1-5-times the IQR. *Descending order by J² statistics value.

(74.5-77.6) in the third, and 74.9% (73.3-76.3) in the fourth quartiles. The pattern was similar for non-Hodgkin lymphoma, but the gap between the first three quartiles and the fourth was much bigger: 59.3% (58.7-60.0) in the first, 57.6% (55.2-58.7) in the second, 55.4% (54.3-56.5) in the third, and 44.7% (43.6-45.8) in the fourth (appendix p 17).

The decrease through quartiles was less uniform for lymphoid malignancies with a poor prognosis: 10-year ASRS for mantle cell lymphoma ranged from $45 \cdot 0\%$ (95% CI $41 \cdot 1-47 \cdot 9$) in the first quartile, to $31 \cdot 3\%$ (28 · 9 – 33 · 8) in the second, $36 \cdot 6\%$ (31 · 5 – 41 · 7) in the third, and 29 · 8% (22 · 3 – 37 · 1) in the fourth quartiles. The

survival pattern for mature T and natural killer cell lymphoma ranged from $48 \cdot 7\%$ (95% CI $46 \cdot 5-50 \cdot 9$) in the first to $45 \cdot 3\%$ ($43 \cdot 3-47 \cdot 3$) in the second, $46 \cdot 0\%$ ($42 \cdot 5-49 \cdot 4$) in the third, and $40 \cdot 9\%$ ($36 \cdot 3-45 \cdot 3$) in the fourth. The corresponding figures for plasma cell neoplasms were $29 \cdot 8\%$ (95% CI $28 \cdot 8-30 \cdot 8$), $29 \cdot 3\%$ ($28 \cdot 5-30 \cdot 1$), $27 \cdot 6\%$ ($26 \cdot 2-28 \cdot 9$), and $22 \cdot 5\%$ ($21 \cdot 2-23 \cdot 8$).

In most countries in the first (high) or second (middlehigh) total national health expenditure quartiles, 10-year ASRS was in line with or above the European mean for common lymphoid malignancies, such as Hodgkin lymphoma, non-Hodgkin lymphoma, follicular lymphoma, and diffuse large B-cell lymphoma (figure 4; appendix pp 8–12).

Among countries in the third (middle-low) total national health expenditure group, 10-year ASRS was on average slightly below or close to the European average for Hodgkin lymphoma and non-Hodgkin lymphoma, respectively (figure 4; appendix pp 8–12).

In most countries in the fourth total national health expenditure quartile, ASRS values were lower than the European mean for many lymphoid malignancies (figure 4; appendix pp 8-12).

The multivariable regression models for each lymphoid malignancy 10-year RER adjusted by age, gender, and total national health expenditure quartiles found that RER increased with age at diagnosis for all lymphoid malignancies (table 3). Multivariable analysis confirmed higher survival for women than for men for all lymphoid malignancies, except precursor B, T, or natural killer cell, or not-otherwise-specified lymphoblastic lymphoma or leukaemia and hairy cell leukaemia.

With reference to the European mean, 10-year RER of all lymphoid malignancies increased with decreasing quartiles of TNHE, and countries in the highest TNHE quartile had lower RER than the European mean. For non-Hodgkin lymphoma, the RER ranged from 0.80 (95% CI 0.79-0.82), to 0.91 (0.90-0.93), 0.94 (0.92-0.96), and 1.45 (1.42-1.48) from the highest to the lowest total national health expenditure quartile.

The generalised linear models re-computed by two-way and three-way interaction effects did not show relevant differences in RERs with respect to the ones that considered only the main effects (appendix pp 21–22). In addition, the analyses performed in the subgroups defined for Hodgkin lymphoma and non-Hodgkin lymphoma did not account for remarkable differences in RERs (appendix p 23), thus showing that the effects of age, gender, and total national health expenditure seemed to be relevant only as main effects.

Discussion

Our population-based study of long-term survival for patients with lymphoid malignancies confirmed the previously reported geographical inequalities in short-tomedium-term survival.⁷ The study also found that



Figure 4: 10-year age-standardised relative survival for Hodgkin lymphoma and non-Hodgkin lymphoma in the EUROCARE-6 countries grouped by total national health expenditure quartile

Data for age-standardised relative survival are presented as percentage, with 95% CI indicated by error bars. The data listed at the top of the figure are the RER estimates for each quartile, with reference to the European mean; the dotted line shows the European mean relative survival. RER=relative excess risk of death. TNHE=total national health expenditure. Graphs for all other lymphoid malignancies are provided in the appendix (pp 8–12).

long-term survival outcomes for patients with lymphoid malignancies are markedly associated with a country's overall investment in the health system (public or private), as measured by per-capita total national health expenditure. The categorisation of total national health expenditure into quartiles, rather than tertiles or quintiles, struck a balance between results granularity, statistical robustness and interpretability, making quartiles a preferred choice for grouping European countries based on their total national health expenditure (appendix pp 24–25). In countries with higher health-care expenditure, 10-year survival was better than in those with lower expenditures. This finding was true for all the lymphoid malignancies, although the survival decrease through the expenditure quartiles was not linear for all of them.

The survival advantage of women compared with men for most lymphoid malignancies, confirmed by multivariable analyses adjusted by age and total national health expenditure and consistent with other studies, might be due to earlier diagnosis, better health behaviours, or lower levels of multimorbidity.³⁰ Life

	Hodgkin Lymphoma	Non-Hodgkin Lymphoma	Chronic Iymphocytic leukaemia/ small Iymphocytic Iymphoma	Lymphoplasmacytic Jymphoma/ Waldenström macroglobulinaemia	Diffuse large B-cell lymphoma	Mantle cell lymphoma	Follicular lymphoma	Marginal zone lymphoma	Mature T and natural killer cell lymphoma	Precursor B-cell, T-cell, natural killer, or not- otherwise- specified lymphoblastic leukaemia	Plasma cell neoplasms	Hairy cell leukaemia
Age groups*												
15-54 years	NA	NA	NA	NA	NA	1 (ref)	NA	1 (ref)	NA	NA	NA	NA
15-44 years	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	NA	1 (ref)	NA	1 (ref)	1 (ref)	1 (ref)	1 (ref)
45-54 years	2·11 (1·87–2·37)	1·42 (1·35–1·49)	1.25 (1.09–1.45)	1.47 (0.89–2.42)	1·64 (1·51–1·79)	NA	1.49 (1.26–1.76)	NA	1·45 (1·27–1·64)	1.96 (1.75–2.19)	1-44 (1-31–1-58)	0.84 (0.38–1.88)
55-64 years	4·20 (3·80-4·64)	2·00 (1·91–2·09)	1.66 (1.45–1.90)	2.03 (1.27–3.24)	2·18 (2·02–2·35)	1.68 (1.43-1.98)	2.46 (2.11–2.86)	2.08 (1.72–2.52)	1·97 (1·76-2·21)	2·71 (2·46–2·99)	1.88 (1.73–2.05)	1.46 (0.69–3.10)
65-74 years	NA	2·92 (2·80–3·05)	2·45 (2·15-2·80)	3·12 (1·97-4·95)	3·11 (2·90–3·35)	2·59 (2·23-3·01)	3·45 (2·97-4·02)	3·18 (2·66–3·81)	2·52 (2·26–2·81)	NA	2.80 (2.58–3.05)	2·17 (1·01-4·68)
≥65 years	11.17 (10.33-12.08)	NA	NA	NA	NA	NA	NA	NA	NA	4.08 (3.76–4.43)	NA	NA
≥75 years	NA	5·54 (5·31–5·78)	4·39 (3·85-5·02)	6.14 (3.88–9.69)	5·64 (5·26–6·05)	4·56 (3·92–5·30)	7.46 (6.40-8.69)	5.66 (4.74–6.78)	3·96 (3·55-4·41)	NA	4.61 (4·24–5·02)	9:31 (4·64-18·68)
Gender												
Men	1.26 (1.18–1.49)	1.16 (1.14–1.19)	1·30 (1·26–1·35)	1.25 (1.13-1.39)	1.09 (1.06–1.12)	1·15 (1·06–1·24)	1·23 (1·15-1·32)	1·31 (1·18–1·46)	1·12 (1·05–1·19)	1.01 (0.94–1.08)	1.02 (1.00–1.05)	1.06 (0.70–1.60)
Women	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Total national health ex	penditure quart	tiles with count	riest									
Low expenditure	1·41 (1·33–1·49)	1·45 (1·42–1·48)	1.66 (1.61–1.71)	1.62 (1.40–1.86)	1.22 (1.17–1.28)	1.26 (1.10-1.43)	1.69 (1.57-1.82)	2·49 (2·27-2·74)	1.26 (1.18–1.35)	1·18 (1·12-1·26)	1.28 (1.25–1.31)	2·15 (1·57–2·93)
Middle-low expenditure	0·94 (0·88–0·99)	0-94 (0-92-0-96)	1.04 (1.01–1.08)	1.09 (0.99–1.20)	0.97 (0.94–1.00)	0.86 (0.79-0.93)	0.85 (0.79–0.90)	0.82 (0.75–0.90)	0-83 (0-78–0-87)	1.05 (0.99–1.11)	0-94 (0-92-0-96)	1-13 (0-80–1-57)
Middle-high expenditure	0·99 (0·94–1·04)	0-91 (0-90-0-93)	0.83 (0.80–0.85)	0.90 (0.83–0.97)	0.96 (0.94-0.99)	1·14 (1·07–1·21)	0-85 (0-81-0-90)	0.70 (0.64–0.76)	1.01 (0.96–1.06)	0.89 (0·84–0·94)	0.94 (0.92-0.95)	0.76 (0.57–1.02)
High expenditure	0.77 (0.72–0.82)	0.80 (0.79–0.82)	0.70 (0.68–0.72)	0.63 (0.57–0.70)	0.88 (0.85–0.90)	0.82 (0.76–0.88)	0-82 (0-77-0-87)	0.70 (0.63–0.77)	0.95 (0.91–1.00)	0.90 (0.85-0.95)	0.89 (0.87–0.90)	0.54 (0.37–0.81)
European mean vs total national health expenditure quartiles	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)	1 (ref)
Data are relative excess risk . Lithuania, Poland, and Slova (UK), and Scotland (UK); hior (UK) and Scotland (UK); hior years for non-Hodgkin Jymp lymphoma, plasma cell neor natural killer cell, or not-oth	of death with 95% ukia; middle-low-e h-expenditure cou homa, chronic lyn blasms, and hairy c erwise-specified ly	s Cl. Distinct mode expenditure counti untries comprise A nphocytic leukaen cell leukaemia; 15- imphoblastic lymp	ls referring to each ries comprise Czec vustria, Denmark, ria/small lympho 54, 55–64, 65–74 yhoma/leukaemia	h lymphoid malignancy we ihia, Italy, Malta, Portugal, S Germany, the Nethelands, ytic lymphoma, lymphopla , and ≥75 years for mantle c †Total national health exp	re adjusted for agi slovenia, Spain, ar Norway, and Swit ssmacyticlympho cell lymphoma an enditure in US\$	e, sex, and total n td Wales (UK); mi Izerland. NA=not ma/Waldenström d marginal zone l ourchasing power	ational health e: ddle-high-exper available. Ref=rc n macroglobulin ymphoma; and : parity per capite	cpenditure. Low-e diture countries c eference group. *C temia, diffuse larg -5-44, 45-54, 55-	xpenditure countr comprise Finland, F sroups for age at d je B-cell lymphoma 64, and ≥65 years	ies comprise Bulga irance, Iceland, Ireli iagnosis were: 15–2 a, follicular lympho for Hodgkin lymph	ria, Croatia, Estoni ınd, England (UK) 4, 45–54, 55–64, ma, mature T and oma and precursc	a, Latvia, Northern Ireland 65-74, and ≥75 natural killer cell or B-cell, T-cell,
Table 3: 10-year relative e	excess risk of dea	ath by age, sex,	and total natior	al health expenditure q	uartile, for each	lymphoid mali	ignancy					

expectancy is better for women than for men everywhere, despite inequality in employment, income, and access to health care, and their increased survival might also be explained by genetic characteristics and immune responses mediated by sex hormones.

The steep relative survival decrease with age, evident for all lymphoid malignancies, indicates that poorer prognostic factors are concentrated in older patients. Older patients are more likely than younger ones to be undertreated because of comorbidity, medical contraindications to full-dose chemotherapy,¹¹ chemotherapyrelated toxicity,¹⁶ and increased risk of undesired effects of anticancer treatments. Social isolation, poverty, and comorbidity are also more frequent among older people and are also associated with delayed cancer diagnoses and poorer outcomes.

The resources that a country can allocate for health care depend on its wealth, and total national health expenditure is directly related to gross national product.⁹ Although the prices of oncological drugs vary across Europe and are lower in some regions than others, countries with lower incomes have more difficulty in allocating resources for expensive diagnostic equipment and treatment modalities, and find it harder to train and employ health personnel in all health sectors, than countries with higher incomes.

The high survival rates for people with Hodgkin lymphoma in countries with high and medium-high national health expenditure indicate that long-standing effective treatments have been deployed well in these countries, whereas the lower survival rates in lower health expenditure countries might be related to a paucity of health-care resources and poor organisation of care.

Clinical trials showed that diffuse large B-cell lymphoma is a potentially curable disease with an overall 60-70% chance of cure with R-CHOP (rituximab plus cvclophosphamide. doxorubicin, vincristine, and prednisone).31 10-year overall survival for patients with diffuse large B-cell lymphoma was 27.6% (95% CI 21.4-34.3) with CHOP only and 43.5% (36.4-50.4) with R-CHOP.³² Other studies showed that overall survival for patients with follicular lymphoma improved significantly since the introduction of rituximab. The Swedish Lymphoma Registry reported 10-year overall survival for patients with follicular lymphoma from 2003 to 2010 to be 92% for those aged 18-49 years, 83% for those aged 50-59 years, 78% for those aged 60-69 years, and 64% for those aged 70 years or older,33 and US and French cohorts' results confirmed improved overall survival for patients with follicular lymphoma with rituximab (10-year overall survival of approximately 80%), compared with without rituximab.34

We suppose that the small differences in survival for patients with diffuse large B-cell lymphoma (and the whole non-Hodgkin lymphoma group) across the second and third total national health expenditure quartiles and among countries of the same quartile might be related to the availability of effective drugs, such as rituximab, and protocols, such as R-CHOP.³⁵ These new options for patients with non-Hodgkin lymphoma have been in clinical use since 2000 and have coincided with many lymphoid malignancy survival gains, especially for patients with follicular lymphoma and diffuse large B-cell lymphoma, as we observed. Unfortunately, we do not have detailed information about each country's adoption of protocols. Notably, new regimens are more expensive than previous ones, for single-agent cost (ie, rituximab or immunotherapy) and for the complexity of treatment schemes.

When prognosis was uniformly poor, as for patients with plasma cell neoplasms or mantle cell lymphoma, there was little difference in 10-year survival and RER through the TNHE quartiles, with differences mainly occurring between the highest and lowest quartiles. RER and survival differences across total national health expenditure quartiles were more important when diagnostic modalities and accuracy differed widely between countries. Chronic lymphocytic leukaemia/small lymphocytic lymphoma is a major diagnostic challenge: the concept of pre-malignant clonal proliferation has been introduced for its classification, with a blood clonal population cutoff of 5000 B lymphocytes per mm^{3.36} Distinction between mantle cell lymphoma, chronic lymphocytic leukaemia, atypical chronic lymphocytic leukaemia, and some marginal zone lymphomas was not straightforward as the differential diagnosis required immunophenotype, and cytogenetic and molecular analyses. Revision of cancer registry procedures to include specific surveillance should help ensure more accurate chronic lymphocytic leukaemia/small lymphocytic lymphoma monitoring. Possible under-registration of lymphoplasmacytic lymphoma/Waldenström macroglobulinaemia and chronic lymphocytic leukaemia/small lymphocytic lymphoma might be due to patients being diagnosed and treated in outpatient clinics, with only more advanced cases or late diagnoses recorded in cancer registry databases.

We found a positive association between total national health expenditure and lymphoid malignancy 10-year survival but not in all countries. The UK—at middle-high national health expenditure quartile—showed lower survival than the European mean for six of the 12 lymphoid malignancies analysed, and Finland had lower survival than the European mean for chronic lymphocytic leukaemia/small lymphocytic lymphoma. In countries with high total national health expenditure, lower survival might be due to inefficient health systems delivering poorer value-for-money care. By contrast, the high survival in Italy for Hodgkin lymphoma and non-Hodgkin lymphoma might indicate good health-care value-for-money and efficient use of scarce resources.

Our study has some limitations. First, this is an observational, retrospective, population-based study with the main goal of providing an understanding of the

state of affairs in lymphoid malignancy survival in Europe and to represent a benchmark for future analyses. The relationship between total national health expenditure and lymphoid malignancy survival could be influenced by various factors, such as diagnostic capacity, treatment accessibility, quality of care, surveillance, and health-care professional expertise. Unfortunately, the only available confounders in this study are age at diagnosis, gender, and period of diagnosis. Including a broader range of confounders would have enhanced the validity of our findings.

Second, this study is based on incidence data referring to 10 years ago on average, but they are the most recent data available.14 Several factors contributed to delays in such a comprehensive study conduction that, notably, ensures maximum geographical representativeness, an essential strength of this study. The lag in cancer registry data recording in Europe, depending on country-specific operational conditions (scarce dedicated resources available, organisation of electronic health-care data, digitisation of processes, and compliance with local privacy regulations) resulted in different data collection times. Moreover, quality controls, which involved multiple revisions and submissions, to ensure maximum result comparability and high data quality, required additional time to permit all data analyses. Plans for the new EUROCARE round include recruitment of patients diagnosed up to 2021 and followed up to the end of 2022. A more flexible strategy to include updates of registry data is envisaged to improve the timeliness of estimations. Furthermore, the process of error checks and correction will be further automated to speed up the analyses.

In addition, changes in the classification of lymphoid malignancies during the study period impaired the conversion of some old codes to newer ones, leading to fairly high proportions of not-otherwise-specified morphologies.

Finally, total national health expenditure might not depict regional cancer registry areas because it represents the whole country. In Italy, most cases were provided by cancer registries in the wealthier northern and central areas. The centralisation of care for haematological malignancies in specialised hospitals can also contribute to relatively good outcomes.³⁷

In conclusion, survival inequalities in patients with lymphoid malignancies uncovered in the past are still evident in more recent years and persist in the long-term follow-up. In countries with low total national health expenditure, policy decisions are needed to increase health-care resources and identify strategies to sustain interventions. Differences in lymphoid malignancy survival among countries with similar total national health expenditure suggest the need for investigation of national health-care policies and health resource utilisation for lymphoid malignancies. In all countries, the implementation and dissemination of efficient, evidence-based models of care, including long-term surveillance, is needed to reduce inequalities. Given the high (and rising) cost of drugs for haematological malignancies, political pressure and negotiations with drug manufacturers to reduce prices of the most expensive drugs should be pursued.

Contributors

MS designed the study, drafted the study protocol, supervised the entire work, and drafted most of the initial version of the report. CV designed the study, drafted the study protocol, and contributed to define quality controls. SR, RL, and RDA did all statistical analyses and wrote the relevant parts of the report. SB prepared all tables and figures, and contributed to statistical analyses and quality controls. MS, CV, SR, SB, ED, and RDA prepared the data and did the quality controls. SR, RL, RDA, MS, CV, SB, and ED accessed and verified the data, MS, CV, RL, SR, SB, RM-G, MM, KI, KP, OV, ED, AB, CDB, SMM, MB, PW, DB, DS, and RDA contributed to interpreting the results. The EUROCARE-6 Working Group revised the study protocol, collected, prepared, and transmitted raw data to the study database, corrected data after quality controls, checked the results of the analyses, and revised the Article. All authors contributed to data interpretation and rewriting of the Article, and reviewed and approved the final version. All the participating cancer registries collected and prepared the data as part of their essential role in cancer control. MS, CV, SR, SB, RL, ED, RDA, AB, and CDB had full access to all the data and all authors had final responsibility to submit for publication.

Declaration of interests

We declare no competing interests.

Data sharing

We analysed pseudonymised data collected from 82 population-based cancer registries, after approval by the Ethics Committee of the National Cancer Institute of Milan (INT73/16; April 21, 2016). We hold these data in trust from each participating registry for the statistical analyses agreed in the EUROCARE-6 protocol. We are not permitted to share individual-level data. Aggregated-level data, in the form of counts, rates, or survival proportions, can be only shared after express permission from the participating registries. These data should be requested by contacting the corresponding author or the Eurocare Secretariat (eurocare. secretariat@istitutotumori.mi.it).

Acknowledgments

This study was partially funded by Italian Ministry of Health "Ricerca Corrente" funds. Funding from the European Commission (Work Programme 2017, grant agreement 801520 HP-JA-2017, "Innovative Partnership for Action Against Cancer") was used to prepare and clean the EUROCARE-6 database. The work of KI and KP was supported by the Estonian Research Council under grant PRG722. We thank Judith Baggot for help with writing of the report in English.

References

- Sung H, Ferlay J, Siegel RL, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021; **71**: 209–49.
- P. Dyba T, Randi G, Bray F, et al. The European cancer burden in 2020: incidence and mortality estimates for 40 countries and 25 major cancers. *Eur J Cancer* 2021; **157**: 308–47.
- 3 Krok-Schoen JL, Fisher JL, Stephens JA, et al. Incidence and survival of hematological cancers among adults ages ≥75 years. *Cancer Med* 2018; 7: 3425–33.
- 4 Usui Y, Ito H, Koyanagi Y, et al. Changing trend in mortality rate of multiple myeloma after introduction of novel agents: a populationbased study. *Int J Cancer* 2020; 147: 3102–09.
- 5 Sterner RC, Sterner RM. CAR-T cell therapy: current limitations and potential strategies. *Blood Cancer J* 2021; 11: 69.
- 5 Chohan KL, Siegler EL, Kenderian SS. CAR-T cell therapy: the efficacy and toxicity balance. *Curr Hematol Malig Rep* 2023; 18: 9–18.
- 7 Sant M, Minicozzi P, Mounier M, et al. Survival for haematological malignancies in Europe between 1997 and 2008 by region and age: results of EUROCARE-5, a population-based study. *Lancet Oncol* 2014; 15: 931–42.

- 8 Burns R, Leal J, Sullivan R, Luengo-Fernandez R. Economic burden of malignant blood disorders across Europe: a population-based cost analysis. *Lancet Haematol* 2016; **3**: e362–70.
- 9 Baili P, Di Salvo F, Marcos-Gragera R, et al. Age and case mixstandardised survival for all cancer patients in Europe 1999–2007: results of EUROCARE-5, a population-based study. *Eur J Cancer* 2015; **51**: 2120–29.
- 10 Dal Maso L, Panato C, Tavilla A, et al. Cancer cure for 32 cancer types: results from the EUROCARE-5 study. *Int J Epidemiol* 2020; 49: 1517–25.
- 11 Moccia AA, Schaff K, Freeman C, et al. Long-term outcomes of R-CEOP show curative potential in patients with DLBCL and a contraindication to anthracyclines. *Blood Adv* 2021; 5: 1483–89.
- 12 Junlén HR, Peterson S, Kimby E, et al. Follicular lymphoma in Sweden: nationwide improved survival in the rituximab era, particularly in elderly women: a Swedish Lymphoma Registry study. *Leukemia* 2015; 29: 668–76.
- 13 Nabhan C, Byrtek M, Rai A, et al. Disease characteristics, treatment patterns, prognosis, outcomes and lymphoma-related mortality in elderly follicular lymphoma in the United States. Br J Haematol 2015; 170: 85–95.
- 14 Botta L, Gatta G, Capocaccia R, et al. Long-term survival and cure fraction estimates for childhood cancer in Europe (EUROCARE-6): results from a population-based study. *Lancet Oncol* 2022; 23: 1525–36.
- 15 Trama A, Vener C, Lasalvia P, Bernasconi A. Late mortality, subsequent malignant neoplasms and hospitalisations in long-term survivors of adolescent and young adult hematological cancers. *Front Oncol* 2022; **12**: 823115.
- 16 Asmis TR, Ding K, Seymour L, et al. Age and comorbidity as independent prognostic factors in the treatment of non-small-cell lung cancer: a review of National Cancer Institute of Canada Clinical Trials Group trials. J Clin Oncol 2008; 26: 54–59.
- 17 Steingrímsson V, Lund SH, Dickman PW, et al. Survival, causes of death, and the prognostic role of comorbidities in chronic lymphocytic leukemia in the pre-ibrutinib era: a population-based study. Eur J Haematol 2022; 108: 145–53.
- 18 Lee SF, Evens AM, Ng AK, Luque-Fernandez MA. Socioeconomic inequalities in treatment and relative survival among patients with diffuse large B-cell lymphoma: a Hong Kong population-based study. *Sci Rep* 2021; 11: 17950.
- 19 Lillini R, Vercelli M, Quaglia A, Micheli A, Capocaccia R. Use of socio-economic factors and healthcare resources to estimate cancer survival in European countries with partial national cancer registration. *Tumori* 2011; 97: 265–74.
- 20 Rossi S, Baili P, Capocaccia R, et al. The EUROCARE-5 study on cancer survival in Europe 1999–2007: database, quality checks and statistical analysis methods. *Eur J Cancer* 2015; **51**: 2104–19.
- 21 Fritz A, Percy C, Jack A, et al. International classification of diseases for oncology (ICD-O), 3rd edn. Geneva, Switzerland: World Health Organization, 2000.
- 22 Swerdlow SH, Campo E, Harris NL, et al. WHO classification of tumours of haematopoietic and lymphoid tissues. WHO classification of tumours, vol 2, revised 4th edition. Lyon, France: IARC Press, 2017.

- 23 Brenner H, Gefeller O, Hakulinen T. Period analysis for 'up-to-date' cancer survival data: theory, empirical evaluation, computational realisation and applications. *Eur J Cancer* 2004; 40: 326–35.
- 24 Pohar Perme M, de Wreede LC, Manevski D. What is relative survival and what is its role in haematology? Best Pract Res Clin Haematol 2023: 36: 101474.
- 25 Corazziari I, Quinn M, Capocaccia R. Standard cancer patient population for age standardising survival ratios. *Eur J Cancer* 2004; 40: 2307–16.
- 26 Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ 2003; 327: 557–60.
- 27 Dickman PW, Coviello E. Estimating and modeling relative survival. *Stata J* 2015; **15**: 186–215.
- 28 Coxe S, West SG, Aiken LS. Generalized linear models. In: Little TD. The Oxford handbook of quantitative methods, vol 2. Oxford: Oxford Library of Psychology, 2013: 26–51.
- 29 Miranda-Galvis M, Tjioe KC, Balas EA, Agrawal G, Cortes JE. Disparities in survival of hematologic malignancies in the context of social determinants of health: a systematic review. *Blood Adv* 2023; 7: 6466–91.
- 30 Dong M, Cioffi G, Wang J, et al. Sex differences in cancer incidence and survival: a pan-cancer analysis. *Cancer Epidemiol Biomarkers Prev* 2020; 29: 1389–97.
- Vodicka P, Klener P, Trneny M. Diffuse large B-cell lymphoma (DLBCL): early patient management and emerging treatment options. OncoTargets Ther 2022; 15: 1481–501.
- 32 Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood* 2010; **116**: 2040–45.
- 33 Junlén HR, Peterson S, Kimby E, et al. Follicular lymphoma in Sweden: nationwide improved survival in the rituximab era, particularly in elderly women: a Swedish Lymphoma Registry study. *Leukemia* 2015; 29: 668–76.
- 34 Sarkozy C, Maurer MJ, Link BK, et al. Cause of death in follicular lymphoma in the first decade of the rituximab era: a pooled analysis of French and US cohorts. J Clin Oncol 2019; 37: 144–52.
- 35 Pulte D, Jansen L, Gondos A, et al. Survival of patients with non-Hodgkin lymphoma in Germany in the early 21st century. *Leuk Lymphoma* 2013; 54: 979–85.
- 66 Call TG, Norman AD, Hanson CA, et al. Incidence of chronic lymphocytic leukemia and high-count monoclonal B-cell lymphocytosis using the 2008 guidelines. *Cancer* 2014; 120: 2000–05.
- 37 Gatta G, Capocaccia R, Botta L, et al. Burden and centralised treatment in Europe of rare tumours: results of RARECAREnet-a population-based study. *Lancet Oncol* 2017; 18: 1022–39.